




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 31/50, 31/445, 31/495, 31/505, 31/535, C07D 211/68, 211/80, 213/02, 239/02, 401/00, 401/04, 471/02, 473/00		A1	(11) International Publication Number: WO 97/05878
			(43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: PCT/US96/12922		(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).	
(22) International Filing Date: 6 August 1996 (06.08.96)			
(30) Priority Data:		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
60/002,094 10 August 1995 (10.08.95) US			
60/014,182 26 March 1996 (26.03.96) US			
9607253.3 4 April 1996 (04.04.96) GB			
9608472.8 25 April 1996 (25.04.96) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(72) Inventors; and		Published	
(75) Inventors/Applicants (for US only): DE LASZLO, Stephen, E. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MANTLO, Nathan, B. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PONTICELLO, Gerald, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). SELNICK, Harold, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). LIVERTON, Nigel, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		With international search report.	

(54) Title: 2,5-SUBSTITUTED ARYL PYRROLES, COMPOSITIONS CONTAINING SUCH COMPOUNDS AND METHODS OF USE



(I)

(57) Abstract

The present invention addresses 2,5-substituted aryl pyrroles of formula (I); or pharmaceutically acceptable salts thereof, as well as compositions containing such compounds and methods of treatment. The compounds are useful for treating Cytokine mediated diseases, which refers to diseases or conditions in which excessive or unregulated production or activity of one or more cytokines occurs. Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8) and Tumor Necrosis Factor (TNF) are cytokines which are involved in immunoregulation and other physiological conditions, such as inflammation. The compounds also have glucagon antagonist activity.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

TITLE OF THE INVENTION

2,5-SUBSTITUTED ARYL PYRROLES , COMPOSITIONS
CONTAINING SUCH COMPOUNDS AND METHODS OF USE

5 CROSS REFERENCE TO RELATED APPLICATIONS

This application is based upon provisional U.S. application no. 60/002,094 filed on August 10, 1995 and upon provisional U. S. application no. 60/014,182 filed on March 23, 1996, priority of which is claimed hereunder.

10

BACKGROUND OF THE INVENTION

The present invention addresses 2,5-substituted aryl pyrroles, as well as compositions containing such compounds and methods of treatment.

15

Cytokine mediated diseases refers to diseases or conditions in which excessive or unregulated production of one or more cytokines occurs. Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are cytokines produced by a variety of cells, which are involved in immunoregulation and other physiological conditions, such as inflammation.

20

IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions. [See, e.g., Dinarello et al., Rev. Infect. Disease, 6, 51 (1984)]. The myriad of known biological activities of IL-1 include the activation of T-helper cells, induction of fever, stimulation of prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase

25

proteins and the suppression of plasma iron levels.

There are many disease states in which IL-1 is implicated. Included among these diseases are rheumatoid arthritis, osteoarthritis, endotoxemia, toxic shock syndrome, other acute or chronic inflammatory diseases, such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis. Recent evidence also links IL-1 activity to diabetes and pancreatic β cells.

35

Excessive or unregulated TNF production has been implicated in mediating or exacerbating rheumatoid arthritis, rheumatoid spondylitis,

- 2 -

osteoarthritis, gouty arthritis, and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS related complex (ARC), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis and pyresis.

Monokines, such as TNF, have been shown to activate HIV replication in monocytes and/or macrophages [See Poli, *et al.*, Proc. Natl. Acad. Sci., 87:782-784 (1990)]. Therefore, inhibition of monokine production or activity aids in limiting HIV progression as stated above for T-cells. TNF has also been implicated in various roles with other viral infections, such as the cytomegalovirus (CMV), influenza virus and the herpes virus.

IL-6 is a cytokine effecting the immune system, hematopoiesis and acute phase reactions. It is produced by several mammalian cell types in response to agents such as IL-1 and is correlated with disease states such as angiofollicular lymphoid hyperplasia.

Interleukin-8 (IL-8) is a chemotactic factor first identified and characterized in 1987. Many different names have been applied to IL-8, such as neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil chemotactic factor (MDNCF), neutrophil activating factor (NAF), and T-cell lymphocyte chemotactic factor. Like IL-1, IL-8 is produced by several cell types, including mononuclear cells, fibroblasts, and endothelial cells. Its production is induced by IL-1, TNF and by lipopolysaccharide (LPS). IL-8 stimulates a number of cellular functions in vitro. It is a chemoattractant for neutrophils, T-lymphocytes and basophils. It induces histamine release from basophils. It causes lysosomal enzyme release and respiratory burst from neutrophils, and it has been shown to increase the surface expression of Mac-1 (CD11b/CD 18) on neutrophils without de novo protein synthesis. There remains a need for treatment, in this field, for compounds which are cytokine suppressive or antagonistic,


- 3 -

i.e., compounds which are capable of inhibiting or antagonizing cytokines such as IL-1, IL-6, IL-8 and TNF.

The compounds of formula I are also useful in treating diseases characterized by excessive IL-8 activity. There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. These diseases include psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury, adult respiratory distress syndrome, thrombosis and glomerulonephritis.


10 SUMMARY OF THE INVENTION

The present invention is directed to a compound represented by formula I:



15

or a pharmaceutically acceptable salt thereof, wherein:



and each independently represent a 5-10 membered aryl or heteroaryl group substituted with R^a groups;

20

wherein a and b represents integers, 0, 1, 2 or 3, such that the sum of a plus b is 1, 2, 3 or 4;



25

represents a heteroaryl group containing from 5 to 10 atoms, 1-4 of which are heteroatoms, 0-4 of which heteroatoms are N and 0-1 of which are O or S, said heteroaryl group being unsubstituted or substituted with 0-3 R^a groups; each R^a independently represents a member

- 4 -

selected from the group consisting of: halo; CN, NO₂, R²¹; OR²³; SR²³; S(O)R²¹; SO₂R²¹; NR²⁰R²³; NR²⁰COR²¹; NR²⁰CO₂R²¹; NR²⁰CONR²⁰R²³; NR²⁰SO₂R²¹; NR²⁰C(NR²⁰)NHR²⁰, CO₂R²³; CONR²⁰R²³; SO₂NR²⁰R²³; SO₂NR²⁰COR²¹; SO₂NR²⁰CONR²⁰R²³; 5 SO₂NR²⁰CO₂R²¹; OCONR²⁰R²³; OCONR²⁰SO₂R²⁰, C(O)OCH₂OC(O)R²⁰; C(NR²⁰)NR²⁰R²³ and CONR²⁰SO₂R²¹;

R¹ is selected from the group consisting of: H, aryl, C₁₋₁₅ alkyl, C₃₋₁₅ alkenyl, C₃₋₁₅ alkynyl and heterocyclyl, said alkyl, aryl, 10 alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: aryl, heteroaryl, heterocyclyl, OR²⁰, SR²⁰, N(R²⁰)₂, S(O)R²¹, SO₂R²¹, SO₂NR²⁰R²³, SO₂NR²⁰COR²¹, SO₂NR²⁰CONR²⁰R²³, NR²⁰COR²¹, NR²⁰CO₂R²¹, NR²⁰CONR²⁰R²³, N(R²⁰)C(NR²⁰)NHR²⁰, CO₂R²⁰, CONR²⁰R²³, 15 CONR²⁰SO₂R²¹, NR²⁰SO₂R²¹, SO₂NR²⁰CO₂R²¹, OCONR²⁰R²³, OCONR²⁰SO₂R²¹; OCONR²⁰R²³ and C(O)OCH₂OC(O)R²⁰;

R² is selected from the group consisting of: H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, halo, NO₂, heterocyclyl, CN, S(O)R²¹, 20 SO₂R²¹, SO₂N(R²⁰)₂, SO₂NR²⁰COR²¹, SO₂NR²⁰CON(R²⁰)₂, COR²⁰, CO₂R²⁰, CONR²⁰R²³, CONR²⁰SO₂R²¹ and SO₂NR²⁰CO₂R²¹, said alkyl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: halo, heterocyclyl, CN, aryl, heteroaryl, R²⁰, OR²⁰, SR²⁰, NR²⁰R²³, S(O)R²¹, SO₂R²¹, 25 SO₂NR²⁰R²³, SO₂NR²⁰COR²¹, SO₂NR²⁰CONR²⁰R²³, NR²⁰COR²¹, NR²⁰CO₂R²¹, NR²⁰CONR²⁰R²³, NR²⁰C(NR²⁰)NHR²⁰, CO₂R²⁰, CONR²⁰R²³, CONR²⁰SO₂R²², NR²⁰SO₂R²¹, SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²¹ and OCONR²⁰R²³;

30 R²⁰ represents a member selected from the group consisting of: H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl, alkenyl, alkynyl, heterocyclyl, aryl and heteroaryl being optionally substituted with 1-3 groups selected from halo, aryl and heteroaryl; 35

- 5 -

R²¹ represents a member selected from the group consisting of: C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl, such alkyl, alkenyl and alkynyl being optionally interrupted with oxo and/or 1-2 heteroatoms selected from O, S, S(O), SO₂ and NR²⁰, said alkyl, 5 alkenyl, alkynyl, aryl and heteroaryl being optionally substituted with from 1-3 of halo, heterocyclyl, aryl, heteroaryl, CN, OR²⁰, O((CH₂)_nO)_mR²⁰, NR²⁰((CH₂)_nO)_mR²⁰ wherein n represents an integer of from 2 to 4, and m represents an integer of from 1 to 3; SR²⁰, N(R²⁰)₂, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, SO₂NR²⁰COR²², SO₂NR²⁰CON(R²⁰)₂, NR²⁰COR²², 10 NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, NR²²C(NR²²)NHR²², CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²²OC(O)R²⁰, C(O)OCH₂OC(O)R²⁰ and OCON(R²⁰)₂;

R²² is selected from the group consisting of: C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl, 15 alkenyl, and alkynyl being optionally substituted with 1-3 halo, aryl or heteroaryl groups;

R²³ is R²¹ or H;

20 R²⁴ is selected from COR²², CO₂R²², CON(R²⁰)₂, SO₂R²² and R²³;

and when two R²⁰ groups are present, when R²⁰ and R²¹ are present, or when R²⁰ and R²³ are present, said two R²⁰ groups, R²⁰ and 25 R²¹ or said R²⁰ and R²³ may be taken in combination with the atoms to which they are attached and any intervening atoms and represent heterocyclyl containing from 5-10 atoms, at least one atom of which is a heteroatom selected from O, S or N, said heterocyclyl optionally containing 1-3 additional N atoms and 0-1 additional O or S atom.

30 Also included in the invention is a pharmaceutical composition which is comprised of a compound of formula I in combination with a pharmaceutically acceptable carrier.

The invention includes a method of treating psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury, 35 adult respiratory distress syndrome, thrombosis and glomerulonephritis, in a mammal in need of such treatment which comprises administering to said

- 6 -

mammal a compound of formula I in an amount which is effective for treating said disease or condition.


Also included in the invention is a method of treating a cytokine mediated disease in a mammal, comprising administering to a mammalian patient in need of such treatment an amount of a compound of formula I which is effective to treat said cytokine mediated disease.

DETAILED DESCRIPTION OF THE INVENTION

The invention is described herein in detail using the terms defined below unless otherwise specified.

The term "alkyl" refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 15 carbon atoms unless otherwise defined. It may be straight, branched or cyclic. Preferred straight or branched alkyl groups include methyl, ethyl, propyl, isopropyl, butyl and t-butyl. Preferred cycloalkyl groups include cyclopentyl and cyclohexyl.

Alkyl also includes a straight or branched alkyl group which contains or is interrupted by a cycloalkylene portion. Examples include the following:



wherein: x plus y = from 0-10 and w plus z = from 0-9.

The alkylene and monovalent alkyl portion(s) of the alkyl group can be attached at any available point of attachment to the cycloalkylene portion.

When substituted alkyl is present, this refers to a straight, branched or cyclic alkyl group as defined above, substituted with 1-3 groups as defined with respect to each variable.

The term "alkenyl" refers to a hydrocarbon radical straight, branched or cyclic containing from 2 to 15 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic (non-resonating) carbon-carbon double bonds may be present. Preferred alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with

- 7 -

respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted when a substituted alkenyl group is provided.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 15 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Preferred alkynyl groups include ethynyl, propynyl and butynyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted when a substituted alkynyl group is provided.

Aryl refers to aromatic rings e.g., phenyl, substituted phenyl and like groups as well as rings which are fused, e.g., naphthyl and the like. Aryl thus contains at least one ring having at least 6 atoms, with up to two such rings being present, containing up to 10 atoms therein, with alternating (resonating) double bonds between adjacent carbon atoms. The preferred aryl groups are phenyl and naphthyl. Aryl groups may likewise be substituted as defined below. Preferred substituted aryls include phenyl and naphthyl substituted with one or two groups.

The term "heteroaryl" refers to a monocyclic aromatic hydrocarbon group having 5 or 6 ring atoms, or a bicyclic aromatic group having 8 to 10 atoms, containing at least one heteroatom, O, S or N, in which a carbon or nitrogen atom is the point of attachment, and in which one additional carbon atom is optionally replaced by a heteroatom selected from O or S, and in which from 1 to 3 additional carbon atoms are optionally replaced by nitrogen heteroatoms. The heteroaryl group is optionally substituted with up to three R^a groups.

Heteroaryl thus includes aromatic and partially aromatic groups which contain one or more heteroatoms. Examples of this type are thiophene, purine, imidazopyridine, pyridine, oxazole, thiazole, pyrazole, tetrazole, imidazole, pyrimidine, pyrazine and triazine.

The groups Ar^1 and Ar^2 represent 5-10 membered aryl or heteroaryl, each of which is substituted with 0 - 3 groups selected from R^a such that a total of 1 to 4 groups is attached to Ar^1 and Ar^2 .

- 8 -

Preferred are phenyl, naphthyl, pyridyl, pyrimidinyl, thiophenyl, furanyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, and isoxazolyl.

The group (HAr) represents a heteroaryl group which contains
5 from 5 to 10 atoms. One to four atoms are heteroatoms which are selected from O, S and N. The heteroaryl group may be unsubstituted or substituted with 0 -3 R^a groups.

Preferred heteroaryl groups represented by (HAr) are as
10 follows: pyridyl, quinolyl, purinyl, imidazolyl, imidazopyridine, and pyrimidinyl.

The terms "heterocycloalkyl" and "heterocyclyl" refer to a cycloalkyl group (nonaromatic) in which one of the carbon atoms in the ring is replaced by a heteroatom selected from O, S, SO, SO₂ or N, and in which
15 up to three additional carbon atoms may be optionally replaced by heteroatoms.

Heterocyclyl is carbon or nitrogen linked; if carbon linked and contains a nitrogen, then the nitrogen may be substituted by R²⁴. Examples of heterocyclyls are piperidinyl, morpholinyl, pyrrolidinyl,
20 tetrahydrofuranyl, tetrahydroimidazo[4,5-c]pyridinyl, imidazolyl, piperazinyl, pyrrolidin-2-onyl, piperidin-2-onyl and the like.

The term "TNF mediated disease or disease state" refers to any and all disease states in which TNF plays a role, either by production of TNF itself, or by TNF causing another monokine to be released, such as but
25 not limited to IL-1 or IL-6. A disease state in which IL-1, for instance, is a major component, and whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF.

The term "cytokine" as used herein means any secreted
30 polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines regardless of which cells produce them. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1),

- 9 -

Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- α) and Tumor Necrosis Factor-beta (TNF- β).

By the term "cytokine interfering or cytokine suppressive amount" is mean an effective amount of a compound of formula I which will, cause a decrease in the *in vivo* levels of the cytokine or its activity to normal or sub-normal levels, when given to the patient for the prophylaxis or therapeutic treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production or activity.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds are contemplated to be within the scope of the present invention.

Throughout the instant application, the following abbreviations are used with the following meanings:

15	Bu	butyl
	Bn	benzyl
	BOC, Boc	t-butyloxycarbonyl
	calc.	calculated
	CBZ, Cbz	Benzyloxycarbonyl
20	CDI	N,N'-carbonyl diimidazole
	FAB-MS	Fast atom bombardment-mass spectroscopy
	HPLC	High pressure liquid chromatography
	KHMDS	Potassium bis(trimethylsilyl)amide
	LAH	Lithium aluminum hydride
25	LHMDS	Lithium bis(trimethylsilyl)amide
	Me	methyl
	MeOH	methanol
	MPLC	Medium pressure liquid chromatography
	NMR	Nuclear Magnetic Resonance
30	Ph	phenyl
	Pr	propyl
	prep.	prepared
	Pyr.	pyridyl
	TMS	Tetramethylsilane

35

- 10 -

One subset of compounds of the invention includes compounds of formula I wherein Ar¹ and Ar² are independently selected from:

- 5 a) phenyl,
 b) pyridyl,
 c) pyrimidinyl,
 d) thiophenyl,
 e) furanyl,
 f) imidazolyl,
10 g) thiazolyl,
 h) isothiazolyl,
 i) oxazolyl,
 j) isoxazolyl, and
 k) naphthyl.

- 15 Within this subset of compounds, all other variables are as previously defined with respect to formula I.

Another subset of compounds of the invention includes compounds of formula I wherein HAr is selected from:

- 20 a) pyridyl,
 b) quinolyl,
 c) purinyl,
 d) imidazolyl,
 e) imidazopyridine, and
 f) pyrimidinyl.

- 25 Within this subset of compounds, all other variables are as originally defined with respect to formula I.

Another subset of compounds of formula I includes compounds wherein R¹ is hydrogen. Within this subset of compounds, all other variables are as originally defined with respect to formula I.

- 30 Another subset of compounds of formula I includes compounds wherein R¹ represents C₁-15 alkyl, unsubstituted or substituted, as originally defined. Within this subset of compounds, all other variables are as originally defined with respect to formula I.

- 35 Another subset of compounds of formula I includes compounds wherein R² represents a member selected from the group consisting of:

- 11 -

- 5
- a) H;
 - b) alkyl;
 - c) halo;
 - d) CN;
 - e) C(O)C₁₋₆ alkyl;
 - f) C(O)C₁₋₆ alkylphenyl;
 - g) CO₂H;
 - h) CO₂C₁₋₆ alkyl
 - i) CO₂C₁₋₆ alkylphenyl;
 - 10 j) CONH₂;
 - k) CONHC₁₋₆ alkyl;
 - l) C(O)N(C₁₋₆ alkyl)₂;
 - m) SO₂NH₂;
 - n) SO₂NHC₁₋₆ alkyl and
 - 15 o) SO₂N(C₁₋₆ alkyl)₂.

Within this subset of compounds, all other variables are as originally defined with respect to formula I.

Preferred compounds of formula I are realized when:

Ar¹ and Ar² are independently selected from:

- 20
- a) phenyl,
 - b) pyridyl,
 - c) pyrimidinyl,
 - d) thiophenyl,
 - e) furanyl,
 - 25 f) imidazolyl,
 - g) thiazolyl,
 - h) isothiazolyl,
 - i) oxazolyl,
 - j) isoxazolyl and
 - 30 k) naphthyl;

one, two or three R^a groups are present and attached to Ar¹ and Ar², and each R^a is independently selected from the group consisting of: halo, R²¹, OR²³, NR²⁰R²³, CO₂R²³, CONR²⁰R²³, SO₂R²¹ and S(O)R²¹, R²⁰, R²¹ and R²³ are as originally defined;

35

HAr is selected from:

- 12 -

- 5
- a) pyridyl,
 - b) quinolyl,
 - c) purinyl,
 - d) imidazolyl,
 - e) imidazopyridinyl and
 - f) pyrimidinyl;

R¹ is: a) H or
b) substituted or unsubstituted alkyl; and

10

R² is selected from the group consisting of:

- 15
- a) H,
 - b) alkyl,
 - c) halo,
 - d) CN,
 - e) C(O)C₁₋₆ alkyl,
 - f) C(O)C₁₋₆ alkylphenyl,
 - g) CO₂H,
 - h) CO₂C₁₋₆ alkyl,
 - 20 i) CO₂C₁₋₆ alkylphenyl,
 - j) CONH₂,
 - k) CONHC₁₋₆ alkyl,
 - l) C(O)N(C₁₋₆ alkyl)₂,
 - m) SO₂NH₂,
 - 25 n) SO₂NHC₁₋₆ alkyl and
 - o) SO₂N(C₁₋₆ alkyl)₂.

A more preferred subset of compounds of formula I is realized when: (R^a)_a-Ar¹ is selected from the group consisting of:

30

- a) phenyl,
- b) 4-fluorophenyl,
- c) 4-chlorophenyl,
- d) 3-fluorophenyl,
- e) 3-chlorophenyl,
- 35 f) thiophen-2-yl,
- g) thiophen-3-yl,

- 13 -

- | | | |
|----|-----|--------------------------|
| | h) | 4-fluorothiophen-2-yl, |
| | i) | 4-fluorothiophen-3-yl, |
| | j) | 5-fluorothiophen-2-yl, |
| | k) | 5-fluorothiophen-3-yl, |
| 5 | l) | 4-chlorothiophen-2-yl, |
| | m) | 4-chlorothiophen-3-yl, |
| | n) | 5-chlorothiophen-2-yl, |
| | o) | 5-chlorothiophen-3-yl, |
| | p) | 3-methylphenyl, |
| 10 | q) | 3,4-dichlorophenyl, |
| | r) | 3-hydroxyphenyl, |
| | s) | 4-hydroxyphenyl, |
| | t) | 3,4-dihydroxyphenyl, |
| | u) | 3-methyl-2-thiophenyl, |
| 15 | v) | 5-methyl-2-thiophenyl, |
| | w) | 4-carboxymethylphenyl, |
| | x) | 3-cyanophenyl, |
| | y) | 4-cyanophenyl, |
| | z) | 2-pyridyl, |
| 20 | aa) | 2-furoyl, |
| | bb) | 3-furoyl, |
| | cc) | 4-methylsulfinylphenyl, |
| | dd) | 4-trifluoromethylphenyl, |
| | ee) | 3-trifluoromethylphenyl, |
| 25 | ff) | 4-methylphenyl, |
| | gg) | 4-t-butoxyphenyl, |
| | hh) | 3,4-dibenzoyloxyphenyl, |
| | ii) | 3-quinolinyl, |
| | jj) | 3-pyridyl, |
| 30 | kk) | 4-pyridyl, |
| | ll) | 2,4-difluorophenyl, |
| | mm) | 3,4-difluorophenyl, |
| | nn) | 4-methylsulfinylphenyl, |
| | oo) | 4-methylsulfonylphenyl, |
| 35 | pp) | 2-methoxyphenyl, |
| | qq) | 3-methoxyphenyl, |

- 14 -

- rr) 4-nitrophenyl,
- ss) 4-aminomethylphenyl, and
- tt) 2-chlorophenyl.

5 (R^a)_b-Ar² is selected from the group consisting of:

- a) 4-(methylthio)-phenyl,
- b) 4-(ethylthio)-phenyl,
- c) 3-(methylthio)-phenyl,
- d) 2-(methylthio)-phenyl,
- 10 e) 3-(ethylthio)-phenyl,
- f) 4-methylsulfonylphenyl,
- g) 4-ethylsulfonylphenyl,
- h) 3-methylsulfonylphenyl,
- i) 2-methylsulfonylphenyl,
- 15 j) 4-methylsulfinylphenyl,
- k) 4-ethylsulfonylphenyl,
- l) 3-methylsulfinylphenyl,
- m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
- n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
- 20 o) 4-methoxyphenyl,
- p) 4-hydroxyphenyl,
- q) 3-methoxyphenyl,
- r) 2-benzyloxyphenyl,
- s) 4-methylthiophen-2-yl,
- 25 t) 4-methylthiophen-3-yl,
- u) 4-acetylamino-phenyl,
- v) 2-pyrimidinyl,
- w) phenyl,
- x) 4-aminomethylphenyl,
- 30 y) 4-cyanophenyl,
- z) 4-fluorophenyl,
- aa) 4-chlorophenyl,
- bb) 4-bromophenyl,
- cc) 4-carboxyethylphenyl,
- 35 dd) 2-fluorophenyl,
- ee) 3-nitrophenyl,

- 15 -

- 5 ff) 4-nitrophenyl,
 gg) 3-fluorophenyl,
 hh) 4-carboxyphenyl,
 ii) 4-aminophenyl,
 jj) 3-aminophenyl,
 kk) 4-(O(CH₂)₃NMe₂)-phenyl,
 ll) 4-(O(CH₂)₂-piperidin-1-yl)-phenyl,
 mm) 2-methoxyphenyl,
 nn) 3-chlorophenyl,
 10 oo) 4-((4-N-COCH₃)piperazin-1-yl)-phenyl,
 pp) 4-trifluoromethylphenyl,
 qq) 4-bromothiophen-2-yl,
 rr) 5-methylthiophen-2-yl,
 ss) 2-benzoxazolyl,
 15 tt) 2-benzofuranyl,
 uu) 2,5-dimethoxyphenyl and
 vv) 4-morpholinylphenyl;

20 with the proviso that when (R^a)_a-Ar¹ represents a), f), g), z), aa), bb), jj) or
 kk), Ar²-(R^a)_b does not represent v), w), ss) or tt);

(R^a)₀₋₃-HAr is selected from the group consisting of:

- 25 a) 4-pyridyl,
 b) 4-(2-methylpyridyl),
 c) 4-(2-aminopyridyl),
 d) 4-(2-methoxypyridyl),
 e) 4-quinolinyl,
 f) 4-pyrimidinyl,
 g) 9-purinyl,
 30 h) 7-(imidazo[4,5-b]pyridinyl),
 i) 4-(3-methylpyridyl),
 j) 2-pyridyl,
 k) 3,5-dimethyl-4-pyridyl,
 l) 3-quinolinyl,
 35 m) 3-pyridazinyl,
 n) 4-(2-aminobenzyl)pyridyl, and

- 16 -

o) 4-(2-amino)pyrimidinyl;

R¹ is: H; and5 R² is selected from the group consisting of:

- a) H,
- b) F,
- c) Cl,
- d) Br,
- 10 e) CN,
- f) C(O)C₁₋₆ alkyl,
- g) C(O)C₁₋₆ alkylphenyl,
- h) CO₂H,
- i) CO₂C₁₋₆ alkyl,
- 15 j) CO₂C₁₋₆ alkylphenyl,
- k) CONH₂,
- l) CONHC₁₋₆ alkyl,
- m) C(O)N(C₁₋₆ alkyl)₂,
- n) SO₂NH₂,
- 20 o) SO₂NHC₁₋₆ alkyl and
- p) SO₂N(C₁₋₆ alkyl)₂.

Another more preferred subset of compounds of formula I is realized when: R^a_a-Ar^l is selected from the group consisting of:

- 25 a) phenyl,
- b) 4-fluorophenyl,
- c) 4-chlorophenyl,
- d) 3-fluorophenyl,
- e) 3-chlorophenyl,
- 30 f) thiophen-2-yl,
- g) thiophen-3-yl,
- h) 4-fluorothiophen-2-yl,
- i) 4-fluorothiophen-3-yl,
- j) 5-fluorothiophen-2-yl,
- 35 k) 5-fluorothiophen-3-yl,
- l) 4-chlorothiophen-2-yl,

- 17 -

- 5
10
15
20
25
30
35
- m) 4-chlorothiophen-3-yl,
 - n) 5-chlorothiophen-2-yl,
 - o) 5-chlorothiophen-3-yl,
 - p) 3-methylphenyl,
 - q) 3,4-dichlorophenyl,
 - r) 3-hydroxyphenyl,
 - s) 4-hydroxyphenyl,
 - t) 3,4-dihydroxyphenyl,
 - u) 3-methyl-2-thiophenyl,
 - v) 5-methyl-2-thiophenyl,
 - w) 4-carboxymethylphenyl,
 - x) 3-cyanophenyl,
 - y) 4-cyanophenyl,
 - z) 2-pyridyl,
 - aa) 2-furoyl,
 - bb) 3-furoyl,
 - cc) 4-methylsulfinylphenyl,
 - dd) 4-trifluoromethylphenyl,
 - ee) 3-trifluoromethylphenyl,
 - ff) 4-methylphenyl,
 - gg) 4-t-butoxyphenyl,
 - hh) 3,4-dibenzyloxyphenyl,
 - ii) 3-quinoliny,
 - jj) 3-pyridyl,
 - kk) 4-pyridyl,
 - ll) 2,4-difluorophenyl,
 - mm) 3,4-difluorophenyl,
 - nn) 4-methylsulfinylphenyl,
 - oo) 4-methylsulfonylphenyl,
 - pp) 2-methoxyphenyl,
 - qq) 3-methoxyphenyl,
 - rr) 4-nitrophenyl,
 - ss) 4-aminomethylphenyl, and
 - tt) 2-chlorophenyl.

(Ra)_b-Ar² is selected from the group consisting of:

- 18 -

- 5 a) 4-(methylthio)-phenyl,
b) 4-(ethylthio)-phenyl,
c) 3-(methylthio)-phenyl,
d) 2-(methylthio)-phenyl,
e) 3-(ethylthio)-phenyl,
f) 4-methylsulfonylphenyl,
g) 4-ethylsulfonylphenyl,
h) 3-methylsulfonylphenyl,
i) 2-methylsulfonylphenyl,
10 j) 4-methylsulfinylphenyl,
k) 4-ethylsulfonylphenyl,
l) 3-methylsulfinylphenyl,
m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
15 o) 4-methoxyphenyl,
p) 4-hydroxyphenyl,
q) 3-methoxyphenyl,
r) 2-benzyloxyphenyl,
s) 4-methylthiophen-2-yl,
20 t) 4-methylthiophen-3-yl,
u) 4-acetylaminophenyl,
v) 2-pyrimidinyl,
w) phenyl,
x) 4-aminomethylphenyl,
25 y) 4-cyanophenyl,
z) 4-fluorophenyl,
aa) 4-chlorophenyl,
bb) 4-bromophenyl,
cc) 4-carboxyethylphenyl,
30 dd) 2-fluorophenyl,
ee) 3-nitrophenyl,
ff) 4-nitrophenyl,
gg) 3-fluorophenyl,
hh) 4-carboxyphenyl,
35 ii) 4-aminophenyl,
jj) 3-aminophenyl,

- 19 -

- 5
10
15
- kk) 4-(O(CH₂)₃NMe₂)-phenyl,
 - ll) 4-(O(CH₂)₂-piperidin-1-yl)-phenyl,
 - mm) 2-methoxyphenyl,
 - nn) 3-chlorophenyl,
 - oo) 4-((4-N-COCH₃)piperazin-1-yl)-phenyl,
 - pp) 4-trifluoromethylphenyl,
 - qq) 4-bromothiophen-2-yl,
 - rr) 5-methylthiophen-2-yl,
 - ss) 2-benzoxazolyl,
 - tt) 2-benzofuranyl,
 - uu) 2,5-dimethoxy-phenyl and
 - vv) 4-morpholinylphenyl;

with the proviso that when (R^a)_a-Ar¹ represents a), f), g), z), aa), bb), jj) or
 15 kk) Ar²-(R^a)_b does not represent v), w), ss) or tt);

(R^a)₀₋₃-HAr is selected from the group consisting of:

- 20
25
30
- a) 4-pyridyl,
 - b) 4-(2-methylpyridyl),
 - c) 4-(2-aminopyridyl),
 - d) 4-(2-methoxypyridyl),
 - e) 4-quinolinyl,
 - f) 4-pyrimidinyl,
 - g) 9-purinyl,
 - h) 7-(imidazo[4,5-b]pyridinyl),
 - i) 4-(3-methylpyridyl),
 - j) 2-pyridyl,
 - k) 3,5-dimethyl-4-pyridyl,
 - l) 3-quinolinyl,
 - m) 3-pyridazinyl,
 - n) 4-(2-aminobenzyl)pyridyl, and
 - o) 4-(2-amino)pyrimidinyl;


35 R¹ is: a) substituted or unsubstituted C₁₋₁₅ alkyl; and

R² is selected from the group consisting of:

- 20 -

- 5
- 10
- 15
- a) H,
 - b) F,
 - c) Cl,
 - d) Br,
 - e) CN;
 - f) C(O)C₁₋₆ alkyl;
 - g) C(O)C₁₋₆ alkylphenyl;
 - h) CO₂H;
 - i) CO₂C₁₋₆ alkyl
 - j) CO₂C₁₋₆ alkylphenyl;
 - k) CONH₂;
 - l) CONHC₁₋₆ alkyl;
 - m) C(O)N(C₁₋₆ alkyl)₂;
 - n) SO₂NH₂;
 - o) SO₂NHC₁₋₆ alkyl and
 - p) SO₂N(C₁₋₆ alkyl)₂.

A further subset of compounds of the invention includes compounds represented by formula I:



wherein:

- 21 -



and



each independently represent a 5-10 membered aryl or heteroaryl group;

5 a and b represents integers, 0, 1, 2 or 3, such that the sum of a plus b is 1, 2, 3 or 4;



represents a heteroaryl group containing from 5 to 10 atoms, 1-3 of which are heteroatoms, 0-3 of which heteroatoms are N and 0-1 of which are O or S, said heteroaryl group being unsubstituted or substituted with 1-3 R^a groups;

10

each R^a independently represents a member selected from the group consisting of: halo; CN, NO₂, R²¹; OR²³; SR²³; S(O)R²¹; SO₂R²¹; NR²⁰R²³; NR²⁰COR²¹; NR²⁰CO₂R²¹; NR²⁰CONR²⁰R²³; NR²⁰SO₂R²¹; NR²⁰C(NR²⁰)NHR²⁰, CO₂R²³; CONR²⁰R²³; SO₂NR²⁰R²³; SO₂NR²⁰COR²¹; SO₂NR²⁰CONR²⁰R²³; SO₂NR²⁰CO₂R²¹; OCONR²⁰R²³; OCONR²⁰SO₂R²⁰, C(O)OCH₂OC(O)R²⁰ and C(NR²⁰)NR²⁰R²³;

20 R¹ is selected from the group consisting of: H, aryl, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, and heterocyclyl, said alkyl, aryl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: aryl, heteroaryl, heterocyclyl, OR²⁰, SR²⁰, N(R²⁰)₂, S(O)R²¹, SO₂R²¹, SO₂N(R²⁰)₂, SO₂NR²⁰COR²¹, SO₂NR²⁰CON(R²⁰)₂, NR²⁰COR²¹, NR²⁰CO₂R²¹, NR²⁰CON(R²⁰)₂, N(R²⁰)C(NR²⁰)NHR²⁰, CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²¹, NR²⁰SO₂R²¹, SO₂NR²⁰CO₂R²¹, OCON(R²⁰)₂, OCONR²⁰SO₂R²¹ and OCONR²⁰R²³;

30 R² is selected from the group consisting of: H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, halo, NO₂ and heterocyclyl, said alkyl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: halo, heterocyclyl, CN, aryl, heteroaryl, R²⁰, OR²⁰, SR²⁰, N(R²⁰)₂, S(O)R²², SO₂R²²,

- 22 -

SO₂N(R²⁰)₂, SO₂NR²⁰COR²², SO₂NR²⁰CON(R²⁰)₂, NR²⁰COR²²,
 NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, NR²²C(NR²²)NHR²², CO₂R²⁰,
 CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²²,
 OCONR²⁰SO₂R²² and OCONR²⁰R²³;

5

R²⁰ represents a member selected from the group consisting of:
 H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and
 heteroaryl, said alkyl, alkenyl and alkynyl being optionally substituted with
 1-3 groups selected from halo, aryl and heteroaryl;

10

R²¹ represents a member selected from the group consisting of:
 C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl,
 said alkyl, alkenyl, alkynyl, aryl and heteroaryl being optionally substituted
 with from 1-3 of halo, heterocyclyl, aryl, heteroaryl, CN, OR²⁰,
 15 O((CH₂)_nO)_mR²⁰, NR²⁰((CH₂)_nO)_mR²⁰ wherein n represents an integer of
 from 2 to 4, and m represents an integer of from 1 to 3; heterocyclyl, SR²⁰,
 N(R²⁰)₂, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, SO₂NR²⁰COR²²,
 SO₂NR²⁰CON(R²⁰)₂, NR²⁰COR²², NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂,
 NR²²C(NR²²)NHR²², CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²²,
 20 NR²⁰SO₂R²², SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²² OC(O)R²⁰,
 C(O)OCH₂OC(O)R²⁰ and OCON(R²⁰)₂;

R²² is selected from the group consisting of: C₁₋₁₅ alkyl, C₂₋₁₅
 15 alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl,
 25 alkenyl, and alkynyl being optionally substituted with 1-3 halo, aryl or
 heteroaryl groups;

R²³ is R²¹ or H;

30 R²⁴ is selected from COR²², CO₂R²², CON(R²⁰)₂, SO₂R²²
 and R²³;

and when two R²⁰ groups are present, when R²⁰ and R²¹ are
 present, or when R²⁰ and R²³ are present, said two R²⁰ groups, R²⁰ and
 R²¹ or said R²⁰ and R²³ may be taken in combination with the atoms to
 35 which they are attached and any intervening atoms and represent
 heterocyclyl containing from 5-10 atoms, at least one atom of which is a

- 23 -

heteroatom selected from O, S or N, said heterocyclyl optionally containing 1-3 additional N atoms and 0-1 additional O or S atom.

Another subset of compounds of the invention includes compounds wherein Ar¹ and Ar² are independently selected from:

- 5 a) phenyl,
- b) pyridyl,
- c) pyrimidinyl,
- d) thiophenyl,
- e) furanyl,
- 10 f) imidazolyl,
- g) thiazolyl,
- h) isothiazolyl,
- i) oxazolyl,
- j) isoxazolyl and
- 15 k) naphthyl;

HAr is selected from the group consisting of::

- a) pyridyl,
- b) quinolyl,
- 20 c) purinyl,
- d) imidazolyl,
- e) imidazopyridine and
- f) pyrimidinyl;

- 25 R¹ is: a) H or
- b) substituted alkyl; and

R² is selected from the group consisting of::

- 30 a) H, or
- b) alkyl and
- c) halo.

Still another subset of compounds of the invention includes compounds wherein Ar¹ is selected from the group consisting of:

- 35 a) phenyl,
- b) 4-fluorophenyl,

- 24 -

- 5
- c) 4-chlorophenyl,
d) 3-fluorophenyl,
e) 3-chlorophenyl,
f) thiophen-2-yl,
g) thiophen-3-yl,
h) 4-fluorothiophen-2-yl,
i) 4-fluorothiophen-3-yl,
j) 5-fluorothiophen-2-yl,
k) 5-fluorothiophen-3-yl,
10 l) 4-chlorothiophen-2-yl,
m) 4-chlorothiophen-3-yl,
n) 5-chlorothiophen-2-yl,
o) 5-chlorothiophen-3-yl,
p) 3-methyl phenyl,
15 q) 3,4 dichlorophenyl and
r) 3-hydroxyphenyl;

(R^a)_b-Ar² is selected from the group consisting of::

- 20 a) 4-(methylthio)-phenyl,
b) 4-(ethylthio)-phenyl,
c) 3-(methylthio)-phenyl,
d) 2-(methylthio)-phenyl,
e) 3-(ethylthio)-phenyl,
f) 4-methylsulfonylphenyl,
25 g) 4-ethylsulfonylphenyl,
h) 3-methylsulfonylphenyl,
i) 2-methylsulfonylphenyl,
j) 4-methylsulfinylphenyl,
k) 4-ethylsulfonylphenyl,
30 l) 3-methylsulfinylphenyl,
m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
o) 4-methoxyphenyl,
p) 4-hydroxyphenyl,
35 q) 3-methoxyphenyl,
r) 2-benzyloxyphenyl,

- 25 -

- s) 4-methylthiophen-2-yl,
- t) 4-methylthiophen-3-yl,
- u) 4-acetylamino-phenyl and
- v) 2-pyrimidinyl;

5

(R^a)₀₋₃-HAr is selected from the group consisting of::

- a) 4-pyridyl,
- b) 4-(2-methylpyridyl),
- c) 4-(2-aminopyridyl),
- 10 d) 4-(2-methoxypyridyl),
- e) 4-quinolinyl,
- f) 4-pyrimidinyl,
- g) 9-purinyl,
- h) 7-(imidazo[4,5-b]pyridinyl), and
- 15 i) 4-(3-methylpyridyl)

R¹ is: H and

R² is selected from the group consisting of:

- 20 a) H,
- b) F,
- c) Cl and
- d) Br.

25 The pharmaceutically acceptable salts of the compounds of formula I include the conventional non-toxic salts or the quarternary ammonium salts of the compounds of formula I formed e.g. from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as

30 hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic and the like.

35 The pharmaceutically acceptable salts of the present invention can be synthesized from the compounds of formula I which

- 26 -

contain a basic or acidic moiety by conventional chemical methods. Generally, the salts are prepared by reacting the free base or acid with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid or base in a suitable solvent or various combinations of solvents.

The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers, including optical isomers, being included in the present invention.

This invention also relates to a method of antagonizing or inhibiting the production or activity of cytokines in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of formula I to antagonize or inhibit cytokine production or activity such that it is regulated down to normal levels, or in some cases to subnormal levels, so as to ameliorate or prevent the disease state.

The compounds of formula I can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of disease states in mammals, which are exacerbated or caused by excessive or unregulated cytokines production, more specifically IL-1, IL-6, IL-8 or TNF, by such mammal's cells, such as but not limited to monocytes and/or macrophages.

Compounds of formula I inhibit cytokines, such as IL-1, IL-6, IL-8 and TNF and are therefore useful for treating inflammatory diseases, such as rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions.

The compounds of formula I may be used to treat other disease states mediated by excessive or unregulated cytokine production or activity. Such diseases include, but are not limited to sepsis, e.g., gram negative sepsis, septic shock, endotoxic shock, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoidosis, bone resorption diseases, such as osteoporosis, reperfusion injury, graft vs. host reaction, allograft rejection, fever and myalgia due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia, secondary to acquired immune deficiency syndrome (AIDS), AIDS and other viral infections, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, pyresis, such as

- 27 -

cytomegalovirus (CMV), influenza virus and the herpes family of viruses such, as Herpes Zoster or Simplex I and II.

The compounds of formula I may also be used in the treatment of inflammation such as for the treatment of rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; inflamed joints, eczema, psoriasis or other inflammatory skin conditions such as sunburn; inflammatory eye conditions including conjunctivitis; pyresis, pain and other conditions associated with inflammation.

The compounds of formula I are normally formulated as pharmaceutical compositions, which are comprised of a compound of formula I and a pharmaceutically acceptable carrier. The compounds of formula I may also be administered in combination with a second therapeutically active compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation.

The pharmaceutical carrier employed may be, for example, solid or liquid. Solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Liquid carriers include syrup, peanut oil, olive oil, water and the like. Similarly, the carrier may include time delay material, such as glyceryl monostearate or glyceryl distearate, alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid dosage form is used, the preparations typically be in the form of a tablet, hard gelatin capsule, a troche or lozenge. The amount of solid will vary widely but preferably will be from about 0.025 mg to about 1 g. When a liquid dosage form is used, the preparation is typically in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid or nonaqueous liquid suspension.

The compounds of formula I may also be administered topically in the form of a liquid, solid or semi-solid. Liquids include solutions, suspensions and emulsions. Solids include powders, poultices and the like. Semi-solids include creams, ointments, gels and the like.

The amount of a compound of formula I, for the methods of use disclosed herein, vary with the compound chosen, the nature and severity of the condition, and other factors. A representative, topical dose

- 28 -

of a compound of formula I is from about 0.01 mg to about 1500 mg, administered one to four, preferably one to two times daily.

While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The active ingredient typically comprises about 0.001% to about 90% w/w.

Drops according to the present invention may comprise sterile aqueous or oil solutions or suspensions, and may be prepared by dissolving the active ingredient in a suitable aqueous solution, optionally including a bactericidal and/or fungicidal agent and/or any other suitable preservative, and optionally including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous liquid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or macrogels. The formulation may

- 29 -


incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as silicas, and other ingredients such as lanolin
5 may also be included.

The methods of the instant invention may also be carried out by delivering the agent parenterally. The term 'parenteral' as used herein includes intravenous, intramuscular, or intradermal and subcutaneous administration. The intravenous and intramuscular forms of administration
10 are preferred. Appropriate dosage forms for such administration may be prepared as described above. The instant invention can also be carried out by delivering the compounds of formula I intranasally, rectally, transdermally or vaginally.

The compounds of formula I may be administered by
15 inhalation. By 'inhalation' is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, include an aerosol formulations and metered dose inhalers.

Compounds of formula I are prepared (see Scheme I) by the reaction of compound 1, or a protected version thereof with an
20 acetophenone (commercially available) in the presence of potassium cyanide followed by treatment with an alkyl or aryl amine, ammonia or equivalent thereof (ammonium acetate) at elevated temperature.

- 30 -

SCHEME I


- 5 Heteroaromatic aldehydes **3** may be converted to their trimethylsilyl cyanohydrins **4**. Deprotonation and reaction with an aldehyde

- 31 -

5 will provide trimethyl silyl protected benzoin **1**. (See, e.g., Hunig, S. et al., *Chem. Ber.* 112, 2062 (1979)).


SCHEME II

5




Condensation of the 1,4-diketone **6** with ammonia or an amine gives rise to pyrroles (Paal Knor Synthesis). Compound **6**, a 1,4-diketone, (see Scheme II) is reacted with ammonia, or a compound that gives rise to ammonia such as ammonium acetate or a primary amine, to provide compounds of formula **1**. This reaction can be conducted in the presence of an acid catalyst, such as acetic acid, or titanium tetrachloride at an elevated temperature. 1,4-diketone **6** is thus regioselectively constructed so that the appropriate groups are present on the pyrrole ring.

- 32 -



- Alkylation of 1-aryl-2-heteroarylethanones **7** with bromoacetophenones or other leaving group substituted acetophenones provides 1,4 diketones **6** (Iyer, R. N.; Gopalachari, R. *Ind. J. Chem.* **11**, 1260, **1973**). Bromoacetophenones are readily prepared by bromination of acetophenones (for example by treatment with bromine in acetic acid or benzyltrimethylammonium bromide).





10

- Ethanone **7** is prepared by the addition of a heteroaryl methyl anion **8** to an activated benzoic acid **9** (for example esters, acid chlorides, nitriles and N-methoxy-N-methyl amides) (see: Wolfe, J. F. et al *J. Org. Chem.* **39**, 2006 **1974** and Kaiser, E. M. et al *Synthesis* 705 **1975** and Ohsawa A. *Chem. Pharm. Bull.* **26**, 3633, **1978**).


15

- 33 -

**7**

- Compound **7** may also be prepared by alkylation of aryl
- 5 trimethyl silyl protected cyanohydrins **10**. Treatment of **10** with lithium diisopropyl amide in THF and addition of a heteroaryl methyl group functionalized with a leaving group L (for example: Br, I, Cl, tosylate, mesylate) followed by acid catalyzed hydrolysis of the silyl cyanohydrin group provides ethanone **7** (Deuchert, K.; Hertenstein, U.; Hunig, S.;
- 10 Wehner, G. Chem. Ber. 112, 2045, 1979).


- 34 -

SCHEME III

5 The reductive cross coupling of a 1,3 diketone **11** with a nitrile **12** in the presence of zinc and titanium tetrachloride also gives rise to compounds of formula I, See Scheme III, (Gao, J. Hu, M.; Chen, J.; Yuan, S.; Chen, W. Tet Lett. 34, 1617, 1993). The 1,3 diketone **11** may be prepared by alkylation of **4** with bromoacetophenones.

10

- 35 -


SCHEME IV

5

The 1,4 diketone **13** can also be prepared as described in Scheme IV. A heteroaryl aldehyde **14** is condensed with a methyl ketone **15** to provide an α,β -unsaturated ketone **16**. In the presence of a catalyst such as cyanide or


- 36 -

a thiazolium salt the aryl aldehyde **17** reacts with **16** to give **13** (Stetter, H. J. et al Heterocyclic Chem. **14**, 573, 1977 and Stetter, H. et. al. Organic Reactions, Vol. 40, 407-496). Condensation of **13** with an amine provides compounds of formula I. Alternatively, variations of Ar^2 may be introduced by addition of Ar^2 aldehydes to alkenes **17b** that are readily available from the ketones **7** described above.


SCHEME V

Intermediate **16** may be prepared by a Horner-Emmons reaction of the anion of **18** with the heteroaryl aldehyde **14**. The reagent **18** may be prepared by reaction of the bromoketone **19** and triethyl phosphite or by reaction of the lithium salt of diethyl methylphosphonate with an ester **21**.

- 37 -


SCHEME VI

- 5 A nitro group may be introduced into the pyrrole nucleus at the 3 position (generic nomenclature - R^2) by electrophilic nitration of a compound such as **22** (or a less advanced intermediate) in the presence of fuming nitric acid and acetic anhydride.
- 10 Halogens may be introduced by electrophilic halogenation with reagents such as XeF_2 ($\text{R}^2 = \text{F}$), N-chlorosuccinimide in DMF ($\text{R}^2 = \text{Cl}$), N-bromosuccinimide in DMF ($\text{R}^2 = \text{Br}$), I_2 in KI ($\text{R}^2 = \text{I}$). Other reagents are available to carry out this conversion, the choice of reagent being dependent on the presence of functional groups that may be sensitive to the reagent being utilized. See e.g., Pyrroles Part 1, R. Alan Jones, ed., Heterocyclic
- 15 Compounds, Vol 48 Part 1, John Wiley, New York, 1990. Pages 348-391.

SCHEME VII

- Introduction of alkyl and heterocyclyl alkyl groups at the 3 position is described. Direct introduction is possible as described in the use of 1,4 diketone **6** as a precursor of compounds of formula **I**. The preparation of the pyrrole **23** containing a hydroxy methyl group at position 3 (R^2) provides an intermediate that is readily elaborated into compounds of formula **I**.
- Acylation of the hydroxyl group with activated acids or isocyanates provides esters and carbamates respectively of formula **I**. Conversion of the hydroxyl group into a leaving group **24** (for example Br, I, Cl, triflate, etc.) enables the introduction of alkyl, heterocyclyl and amines and thiol groups by displacement with a nucleophile. Suitable nucleophiles include for example, an alkyl or heterocyclyl anion, a primary or secondary amine or a thiol.

- 39 -

SCHEME VIII


5

Hydroxy methyl substituted pyrroles **23** may be prepared by reduction of esters **25a** through the reaction with a reducing agent such as

lithium aluminum hydride. The ester **25a** may be prepared by treatment of 1,2-disubstituted-2 halo ketones **22** with 3-keto esters **27** and ammonia or an amine, producing ester **25a** (Hantzsch. Ber. Dtsch. Chem. Ges. **23**, 1474, **1890**). Alternatively, a 2-amino ketone **24** reacts with a 3-keto ester **27** to produce **25**.

A further method of synthesis of **23** is via reduction of the aldehyde **26** with a reducing agent such as sodium borohydride. The aldehyde may be prepared by treatment of the R³ -unsubstituted pyrrole **25** with the Vilsmeier reagent (POCl₃/DMF).

SCHEME IX




The ester and nitrile of formula I may be prepared as shown in Scheme IX by treatment of halo ketones **22** with keto esters or keto nitriles **23a** with ammonia or an amine producing ester **I** (Hantzsch. Ber. Dtsch. Chem. Ges. **23**, 1474, **1890**). Alternatively a 2-amino ketone **24** reacts with a 3-keto ester **23a** to produce **I**. A further method of synthesis of


- 41 -

Compounds of formula I is by oxidation and esterification of aldehyde 26. The aldehyde is prepared by treatment of the pyrrole 25 with the Villsmeyer reagent (POCl_3/DMF).

5

SCHEME X

- 42 -




- The pyrrole **22** prepared as described herein may be silylated on the nitrogen atom to give **27** by treatment with a silyl chloride and base in a solvent such as methylene chloride. The pyrrole **27** may then be sulphenylated with a sulphenylchloride under basic conditions to provide **28** (J. Org. Chem. 6317 1990). Oxidation of **28** with a reagent such as m-chloroperoxybenzoic acid will give the sulphone **29**. Removal of the silyl group and derivatization of the pyrrole will give compounds of Formula I.
- Compound **22** may also be converted to the sulphide **30** by reaction of **22** with a symmetrical sulfoxide in the presence of trimethylsilylchloride (TMSCl) to give **30**. Oxidation of **30** with a reagent such as m-chloroperoxybenzoic acid will give **29**. The silyl pyrrole **27** may also be acylated with an acid chloride to give the ketone **31**.

- 43 -

- Removal of the silyl group from **29** and **31** and derivatization of the pyrrole will give compounds of formula I. Pyrroles such as **22** may also be sulfinylated directly without N-protection, by treatment with sulphinyl chloride in a solvent such as dichloromethane at 0°C (J. Org. Chem. 5336, 1980). Oxidation as described above thus provides pyrroles of formula I where R³ is SO₂R²¹.

SCHEME XI




10

The amino acid ester **32** may be acylated with an acid **33** that is suitably activated (acid chloride or other activating group used in amide coupling reactions) to give **34** (Scheme XI). Hydrolysis of the ester protecting group will provide **35**. Cyclization by treatment with an acid

- 44 -

activating group such as dicyclohexylcarbodiimide (DCC) will give the oxazolium species **36**. Addition of an alkyne **37** to **36** may give a pyrrole of Formula I via a 3+2 cycloaddition followed by loss of carbon dioxide. Various R^3 groups may be incorporated in this manner.

5

SCHEME XII

10 Aryl and heteroaryl rings may also be appended to the pyrrole ring system by utilization of organometallic coupling technology (Kalinin, V. Synthesis 413 **1991**). See Scheme XIII. The pyrrole ring may function as an electrophile or as a nucleophile.

15 Any of the three appended aromatic or heteroaromatic rings may be attached to the pyrrole ring system (Alvarez, A. J. et al. J. Org. Chem. 1653, **1992** (use of boronic acid and tributyl stannanes for coupling to aromatic and heteroaromatic rings)). Attachment of pyrrole pendant groups may be carried out with or without other Ar, HAr, R^2 or R^3 groups attached.

- 45 -

The synthesis of pyrroles containing nucleophilic groups for coupling reactions depends on the pyrrole substitution pattern. Lithium anions are prepared by metalation of a regioselectively halogenated pyrrole, or the regioselective deprotonation of the pyrrole preferably by the use of a directing functional group. The resulting anion may then be trapped by a trialkyl stannyl halide or a trialkyl borate or transmetalated to magnesium or zinc by treatment with appropriate halide salts. A further method used to incorporate a trialkyl stannyl group is the coupling of a bromo, iodo or triflate substituted pyrrole with hexalkylditin in the presence of a palladium catalyst.

The synthesis of pyrroles incorporating electrophilic groups may be carried out by the regioselective halogenation of a pyrrole (Pyrroles, Part 1, R. Alan Jones, ed., Heterocyclic Compounds Vol 48 Part 1, John Wiley, New York, 349-391, 1990). The regioselectivity of halogenation will depend on the size, nature and substitution position on the pyrrole ring as well as the presence or absence of the N-alkyl protecting group. Triflates may be prepared by acylation of hydroxy pyrroles with triflic anhydride.

The reaction conditions used will depend on the nature of the coupling species. In the case of magnesium, zinc and stannyl coupling reactions the solvent employed is toluene, or DMF under anhydrous conditions. In the case of boronic acid couplings, a heterogenous system is used of water, toluene, and dimethoxyethane, or ethanol in the presence of a base such as sodium carbonate, or bicarbonate. In general, the reaction takes place at an elevated temperature (80-100 °C.). The catalysts used will most likely depend on the structure of the components to be coupled as well as the functional groups and belong to the group consisting of tetrakis(triphenyl)phosphinepalladium (0), or palladium bis triphenyl phosphine dichloride.

Coupling of alkenes or alkynes with 4-halo pyrroles (Heck reaction, see Kalinin, V. Synthesis 413 1991 for a review) will give rise to R² (generic nomenclature) alkenyl and alkynyl substituted pyrroles that may be reduced or otherwise modified to provide compounds of formula I.

Functional groups such as halogens, sulfides, nitro groups, ethers and other groups stable to the reaction conditions used in the linear synthesis of the pyrroles are incorporated in the initial steps of the reaction sequence. Sulfides may be oxidized to sulfoxides and sulfones with

- 46 -

reagents such as m-chloroperbenzoic acid. Sulfides may also be converted to sulfonyl chlorides by oxidation and chlorination by chlorine in water.

Primary amines are prepared from nitro groups by catalytic (Pd/C, H₂ or Raney Nickel, H₂) or chemical means (CoCl₂, NaBH₄).

- 5 Alkylation of amines to give secondary and tertiary amines is achieved by reductive alkylation (aldehyde, NaCNBH₄) or alkylation with an alkyl group substituted with a leaving group in the presence of a base such as K₂CO₃. Tertiary amines may, alternatively, be carried through the reaction sequence to the pyrroles. Acylation of primary or secondary amines with
10 activated acids, chloroformates, isocyanates and chlorosulfonates will give rise to amides, carbamates, ureas and sulfonamides, respectively.

- Other methods of preparing amides and ureas are useful; such as for example, treatment of the amine with phosgene, or an equivalent thereof, followed by acylation of an alcohol or amine with the intermediate
15 activated chloroformamide.

- Carboxylic acids are best introduced as esters early in the synthesis. Saponification will provide carboxylic acids. Transesterification or esterification of the acids will give esters. Carboxylic acids may be converted to amides by activation and reaction with amines. Phenols are
20 best introduced in a protected form early in the synthetic sequence to the pyrrole. Removal of the protecting group provides a phenol which may subsequently be alkylated in the presence of an alkylating agent and base to give an ether, or acylated with an isocyanate to give carbamates. Phenols may be converted to aryl ethers by reaction with an aryl bismethane in the
25 presence of copper II acetate.

Aryl and heteroaryl groups may be attached to pyrrole pendant aryl and heteroaryl groups by application of coupling chemistry technology as outlined above.

- The sequence and conditions of the reaction steps is dependent
30 on the structure and functional groups present. Protecting groups may be necessary and may be chosen with reference to Greene, T.W., et al., Protective Groups in Organic Synthesis, John Wiley & Sons, Inc., 1991. The blocking groups are readily removable, i.e., they can be removed, if
35 desired, by procedures which will not cause cleavage or other disruption of the remaining portions of the molecule. Such procedures include chemical and enzymatic hydrolysis, treatment with chemical reducing or oxidizing

- 47 -

agents under mild conditions, treatment with fluoride ion, treatment with a transition metal catalyst and a nucleophile, and catalytic hydrogenation.

Examples of suitable hydroxyl protecting groups are: t-butylmethoxyphenylsilyl, t-butoxydiphenylsilyl, trimethylsilyl, triethylsilyl, 5 o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, benzyloxycarbonyl, t-butylloxycarbonyl, 2,2,2-trichloroethyloxycarbonyl, and allyloxycarbonyl. Examples of suitable carboxyl protecting groups are benzhydryl, o-nitrobenzyl, p-nitrobenzyl, 2-naphthylmethyl, allyl, 2-chloroallyl, benzyl, 2,2,2-trichloroethyl, trimethylsilyl, t-butyl dimethoysilyl, t- 10 butyldiphenylsilyl, 2-(trimethylsilyl)ethyl, phenacyl, p-methoxybenzyl, acetonyl, p-methoxyphenyl, 4-pyridylmethyl and t-butyl.

The following examples are illustrative and are not limiting of the compounds of this invention.

15 PREPARATIVE EXAMPLE 1

4-Fluoro-2-(4-pyridyl)acetophenone (1)

To a solution of lithium diisopropylamide (Aldrich Chemical Co. 2.0 M in heptane, THF and ethylbenzene) 3.1 mL (6.3 mmol) in 6 mL of anhydrous THF at -78°C under nitrogen was added 0.5 g (5.3 mmol) of 4- 20 picoline dropwise. The reaction mixture was stirred for 20 minutes and then treated with a solution of 0.9 g (5.3 mmol) of 4-fluoro-(N-methyl-N-methoxy)-benzamide in THF. The reaction mixture was warmed to 0°C and quenched by addition of 10 mL of brine. The mixture was extracted with ethyl acetate (3 x 10 mL) and the combined organic phases were dried over 25 MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo to give an orange solid (4-Fluoro-2-(4-pyridyl)acetophenone). ¹H NMR (CDCl₃ 300 MHz): 4.23 s (d, 2H), 7.1-7.18 m (4H), 8.02 (dd, 2H), 8.55 (dd, 2H).

30 PREPARATIVE EXAMPLE 2


4-Fluoro-2-(2-pyridyl)acetophenone (2)

To a solution of lithium diisopropylamide (Aldrich Chemical Co. 2.0M in heptane, THF ethylbenzene) 5.2 mL (10.5 mmol) in 6 mL of anhydrous THF at -78°C under nitrogen was added 0.93 g (10 mmol) of 2- 35 picoline dropwise. The reaction mixture was stirred for 20 minutes and then treated with a solution of 1.71 g (5.3 mmol) of 4-fluoro-(N-methyl-N-

- 48 -

methoxy)-benzamide in THF. The reaction mixture was warmed to 0°C and quenched by addition of 10 mL of brine. The mixture was extracted with ethyl acetate (3 x 10 mL) and the combined organic phases were dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo to give a solid. H¹ NMR (CDCl₃, 300 MHz): 4.49 (s); 6.0 (s); 6.97 (m); 7.-3-7.12 (m); 7.62 (m); 7.82 (m); 8.10 (dd); 8.28 (bd); 8.57 (bd). The compound exists in a keto/enol equilibrium as determined by H¹-NMR. FAB ms:216 (M⁺+1).

- 49 -


PREPARATIVE EXAMPLES 3-10

- 5 The following compounds are prepared in the manner described above:

TABLE I		
$ (\text{R}^a)_a - \text{Ar}^I - \text{C}(=\text{O}) - \text{CH}_2 - \text{HAr} - (\text{R}^a)_{0-3} $		
Prep Ex #	$(\text{R}^a)_a - \text{Ar}^I$	$\text{HAr} - (\text{R}^a)_{0-3}$
3	phenyl	4-pyridyl
4	phenyl-4-F	4-(3-methyl)-pyridyl
5	phenyl-4-F	4-quinolinyl
6	phenyl-3-Cl	4-pyridyl
7	phenyl-2-Cl	4-pyridyl
8	phenyl-4-F	4-pyrimidinyl
9	phenyl-4-F	4-(2-Me)-pyridyl*
10	phenyl-3,4-di-F	4-pyridyl

- 10 *Purified by chromatography from the side product formed from alkylation of the 2-methyl group of 2,4-dimethylpyridine.

- 50 -

PREPARATIVE EXAMPLE 11 (Method 1)


5

To a solution of 0.14 g (0.67 mmol) of 4-fluoro-2-(4-pyridyl)acetophenone (1) from Preparative Example 1 in 2 mL of anhydrous DMSO under nitrogen at room temperature was added 0.67 mL of a 1.0M solution of sodium hexamethyl disilazide in THF. After 15 minutes a solution of 4-fluoro-bromoacetophenone 0.14 g (0.67 mmol) in DMSO was added dropwise. The reaction mixture was diluted with 5 mL of water after 30 minutes and extracted with ethyl acetate (3x 10 mL). The combined organic extracts were washed with brine and dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo to give an oil. ¹H NMR (CDCl₃ 300 MHz): 3.26 (dd, 1H); 4.12 (dd, 1H); 5.23 (dd, 1H); ; 7.04-7.13 (m, 4H); 7.27 (dd, 2H); 7.95-8.05 (m, 4H); 8.51 (d, 2H).

15

PREPARATIVE EXAMPLE 12 (Method 2)

20



0.2 g (1.02 mM) of the product of Preparative Example 3 in 3 mL of dry DMF under nitrogen was treated with 48.7 mg of 60 % sodium hydride dispersion in oil. The reaction mixture was stirred at room temperature for one hour. 298 mg (1.22 mM) of 2-bromo-4'-nitroacetophenone dissolved in 1 mL of DMF was added dropwise. The

25

- 51 -

- reaction mixture was stirred at room temperature for 2 hours and was then treated with water and 10% citric acid solution. Ethyl acetate was added and the layers were separated and the aqueous phase was extracted with ethyl acetate. the organic phase was dried over MgSO_4 , filtered and was concentrated in vacuo. The residue was purified by flash chromatography over silica gel eluting with 25-50% EtOAc/Hexanes to give the desired product.

PREPARATIVE EXAMPLES 13-37

- 10 The compounds of preparative examples 13-37 were prepared using starting materials from the preparative examples above.






TABLE II



Prep. Example #	$(R^a)_a$ -Ar ¹	Ar ² -(R ^a) _b	HA ^r -(R ^a) ₀₋₃	Method
13	Ph-4-F	Ph-4-OMe	4-Pyridyl	1
14	Ph-4-F	Ph-2,5-di-OMe	4-Pyridyl	1
15	Ph-4-F	Ph-4-Br	4-Pyridyl	1
16	Ph-4-F	Ph-4-Cl	4-Pyridyl	1
17	Ph-4-F	Ph-4-SMe	4-Pyridyl	1
18	Ph-4-F	Ph-4-OMe	2-Pyridyl	1
19	Ph-4-F	Ph-4-Br	2-Pyridyl	1
20	Ph-4-F	Ph-4-Cl	2-Pyridyl	1
21	Ph-4-F	Ph-2,5-di-OMe	2-Pyridyl	1
22	Ph	Ph-4-OMe	4-Pyridyl	1
23	Ph	Ph-4-Cl	4-Pyridyl	1
24	Ph	Ph-2,5-di-OMe	4-Pyridyl	1
25	Ph	Ph-4-F	4-Pyridyl	1
26	Ph	Ph-4-COOEt	4-Pyridyl	2
27	Ph	Ph-2-F	4-Pyridyl	2
28	Ph	Ph-3-NO ₂	4-Pyridyl	1
29	Ph-4-F	Ph-4-SMe	4-(3-Me)-pyridyl	1
30	Ph	Ph-3-F	4-Pyridyl	2
31	Ph	Ph-2-NO ₂	4-Pyridyl	2
32	Ph-4-F	Ph-4-SMe	4-quinolinyl	1
33	Ph-2-Cl	Phenyl	4-Pyridyl	1

- 53 -


34	Ph-3-Cl	Ph-4-SMe	4-Pyridyl	1
35	Ph-4-F	Phenyl	4-pyrimidinyl	1
36	Ph-4-F	Ph-4-SMe	4-(2-Me)-pyridyl	1
37	Ph-3,4-F	Ph-4-SMe	4-Pyridyl	1

PREPARATIVE EXAMPLE 38

5 To 4.4 mL of dry pyridine under nitrogen was added 2.14 g (0.02 mol) of pyridine-4-carboxaldehyde followed by 2.4 g (0.02 mol) acetophenone and 1.46 g (0.02 mol) diethylamine. The solution was refluxed for 2.5 hours, cooled to room temperature and poured into 100 mL of ice water containing 10 mL of concentrated hydrochloric acid. The

10 resulting solution was adjusted to pH 5.0 by addition of 1N NaOH solution while stirring rapidly. The mixture was filtered and the residue was washed with 15 mL of water. The solid was dried in vacuo to give the product.

^1H NMR (CDCl_3 300 MHz): 7.42-7.56 (m, 3H); 7.59-7.65 (m, 1H); 7.68 (d, 2H); 8.11 (dt, 2H); 8.69 (bd, 2H).

PREPARATIVE EXAMPLE 391-(4-chlorophenyl)-4-phenyl-2-(4-pyridyl)-butan-1,4-dione

20 To 0.019 g (0.039 mMol) of sodium cyanide in 2 ml of dry DMF at 30°C was added a solution of 4-chlorobenzaldehyde in 1.5 mL of DMF over 20 minutes. A thick slurry formed. After 30 minutes a solution of the product of preparative example 38 in 1.5 mL of DMF was added

- 54 -

dropwise. The mixture was agitated and stirred for 3 hours. The mixture was diluted with 30 ml of water and extracted with ethyl acetate (2 x 15 mL). The organic phase was washed with brine (1 x 15 mL) and dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo to give the title compound as an oil.


PREPARATIVE EXAMPLE 40

4-phenyl-2-(4-pyridyl)-1-(4-trifluoromethylphenyl)-butan-1,4-dione

A solution of 0.15 g (0.71 mMol) of 1-phenyl-3-(4-pyridyl)-ethene-1-one (Preparative Example 38), 0.14 g (7.8 mMol) of 4-trifluoromethylbenzaldehyde, 0.035 g (0.35 mMol) of triethylamine and 20 mg (0.07 mMol) of 3,4-dimethyl-5-(2-hydroxyethyl)-thiazolium iodide in 3 mL on ethanol was heated at 80°C for 3 hours. The reaction mixture was cooled to room temperature, diluted with 10 mL of water and was washed with water and brine and dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo and the residue was purified by MPLC over silica gel eluting with 2.5% MeOH/CH₂Cl₂ to give the desired product.

¹H NMR (CDCl₃ 300 MHz): 7.42-7.56 (m, 3H); 7.59-7.65 (m, 1H); 7.68 (d, 2H); 8.11 (dt, 2H); 8.69 (bd, 2H).

- 55 -

PREPARATIVE EXAMPLES 41-60

- 5 Following the procedure described above the following compounds are prepared:

TABLE III	
Prep Ex.#	(R ^a) _a —Ar ¹
41	4-methylthio-Ph
42	4-Me-Ph
43	4-t-BuO-Ph
44	3,4-dichloro-Ph
45	3,4-dibenzoyloxy-Ph
46	2-thiophenyl
47	3-furoyl
48	3-Cl-Ph
49	2-pyridyl
50	4-CN-Ph
51	4-methoxycarbonyl-Ph


- 56 -

52	5-Me-2-thiophenyl
53	3-Me-2-thiophenyl
54	3-quinoliny
55	3-Pyridyl
56	4-Pyridyl
57	4-F-Ph
58	2,4-di-F-Ph
59	3-CN-Ph
60	3,4-di-F-Ph

Ph = phenyl; Pyr = pyridyl; 3,4-di-F = 3, 4-difluoro .


PREPARATIVE EXAMPLE 61

5



To a 2 liter 3 neck flask equipped with a mechanical stirrer under N₂ was added 54.6 g (0.59 m) diisopropylethylamine and 150 mL of THF. The solution was cooled to -20°C and treated with 268 ml (0.67 m) of 2.5 M butyl lithium over 20 minutes. To the reaction mixture was added 125 g (0.56 mMol) of 4-(t-butyl dimethylsilyloxymethyl)pyridine in 100 ml of THF over 30 minutes. The reaction mixture was stirred for 1 hour at -15°C and then treated with a solution of 108 g (0.59 mMol) of 4-fluoro-(N-methyl-N-methoxy)-benzamide dissolved in 100 mL of THF dropwise. The reaction was warmed to 0°C and stirred for 1 hour and then was warmed to room temperature and was quenched by addition of 1 liter of 20% NH₄Cl solution. The aqueous phase was extracted with EtOAc (3 x 500 mL). The combined organic phases were washed with water (1 x 500 mL), 1 x 500 mL brine and were dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo to give a dark oil. The product was purified by flash chromatography over silica gel eluting with 10-20% EtOAc/hexanes.

- 57 -

EXAMPLE 12-(4-fluorophenyl)-5-(4-fluorophenyl)-3-(4-pyridyl)-pyrrole

5

Method 1: To a solution of 0.35 g (0.99 mmol) of the compound of preparative example 11 in 15 mL of glacial acetic acid was added 0.35 g (4.7 mmol) ammonium acetate. The mixture was heated to 90-110°C over 10 hours at which time a further 1 g of ammonium acetate was added. Heating was continued at 110°C for 6 hours. The reaction mixture was concentrated to 50% of the original volume, and 25 mL of water was gradually added. A solid formed, which was filtered and dried in vacuo to give the title compound. ^1H NMR (CDCl_3 300 MHz): 6.61 (s, 1H); 6.98-7.05 (m, 4H); 7.20 (dd, 2H); 7.33 (dd, 2H); 7.53 (dd, 2H); 8.29 (d, 2H). FAB ms: 333 (M^++1).


Method 2: The condensation in method 1 is followed by an alternative work-up procedure. The reaction mixture was diluted with 5 mL of water and extracted with ethyl acetate (3 x 4 mL). The organic extracts are dried over MgSO_4 , filtered and the filtrate concentrated in vacuo. The residue was purified by rotary chromatography to give the desired product.

EXAMPLES 2-43

0.15-0.2 g of the 1,4-diketone from Preparative Examples 13-37 and 39-60 is dissolved in 3 mL of acetic acid to which is added 1.0 g of ammonium acetate. The mixture is heated at 110°C for 1.5-10 hours. The work-up of Method 1 or 2 is then utilized to isolate the compounds (Examples 2-50) listed below in Table IV.

TABLE IV

- 58 -




Ex #	$(R^a)_a$ -Ar ¹	Ar ² -(R ^a) _b	$(R^a)_{0-3}$ -HAr	FAB ms
2	Ph-4-F	Ph-4-OMe	4-Pyridyl	345
3	Ph-4-F	Ph-2,5-di-OMe	4-Pyridyl	375
4	Ph-4-F	Ph-4-Br	4-Pyridyl	393/395
5	Ph-4-F	Ph-4-Cl	4-Pyridyl	349
6	Ph-4-F	Ph-4-OMe	2-Pyridyl	345
7	Ph-4-F	Ph-4-Br	2-Pyridyl	393/395
8	Ph-4-F	Ph-4-Cl	2-Pyridyl	349
9	Ph-4-F	Ph-2,5-di-OMe	2-Pyridyl	375
10	Ph	Ph-4-OMe	4-Pyridyl	327
11	Ph	Ph-4-Cl	4-Pyridyl	331
12	Ph	Ph-2,5-di-OMe	4-Pyridyl	357
13	Ph	Ph-4-F	4-Pyridyl	315
14	Ph-4-Cl	Ph	4-Pyridyl	331
15	Ph-4-F	Ph-4-SMe	4-Pyridyl	361
16	Ph-4-SMe	Ph	4-Pyridyl	325
17	4-CF ₃ -Ph	Ph	4-Pyridyl	365
18	4-Me-Ph	Ph	4-Pyridyl	311
19	4-tBuO-Ph	Ph	4-Pyridyl	369
20	3,4-Cl-Ph	Ph	4-Pyridyl	366
21	3,4-di-(OBn)-Ph	Ph	4-Pyridyl	509
22	3-Cl-Ph	Ph	4-pyridyl	331
23	4-CN-Ph	Ph	4-Pyridyl	322

- 59 -

24	4-(COOMe)-Ph	Ph	4-Pyridyl	355
25	Ph	4-CO ₂ Et-Ph	4-Pyridyl	369
26	Ph	2-F-Ph	4-Pyridyl	315
27	Ph	3-NO ₂ -Ph	4-Pyridyl	342
28	5-Me-thiophen-2-yl	Ph	4-Pyridyl	317
29	3-Me-thiophen-2-yl	Ph	4-Pyridyl	317
30	4-F-Ph	4-SMe-Ph	4-(3-Me)-Pyridyl	375
31	Ph	2-NO ₂ -Ph	4-Pyridyl	342
32	Ph	3-F-Ph	4-Pyridyl	315
33	Ph	4-NO ₂ -Ph	4-Pyridyl	342
34	4-F-Ph	Ph	4-Pyridyl	315
35	2,4-F-Ph	Ph	4-Pyridyl	333
36	3-CN-Ph	Ph	4-Pyridyl	322
37	3,4-F-Ph	Ph	4-Pyridyl	333
38	4-F-Ph	4-(SMe)-Ph	4-quinolinyl	411
39	2-Cl-Ph	Ph	4-pyridyl	331
40	3-Cl-Ph	4-(SMe)-Ph	4-pyridyl	377
41	4-F-Ph	Ph	4-pyrimidinyl	316
42	4-F-Ph	4-(SMe)-Ph	4-(2-methyl)-pyridyl	375
43	3,4-di-F-Ph	4-(SMe)-Ph	4-pyridyl	379

- 60 -

EXAMPLE 442-(4-fluorophenyl)-5-(4-methylsulfinylphenyl)-3-(4-pyridyl) pyrrole

- 5 To a solution of Example 15 (55.5 mg (0.15 mmol) in 2 ml of acetic acid and 1.4 ml of water was added potassium persulfate (50.0 mg (0.18 mmol). After stirring for 1.5 hours at room temperature the solution was neutralized by addition of ammonium hydroxide solution. The solid product was recovered by filtration and purified by flash chromatography
- 10 eluting with 5% MeOH/CH₂Cl₂ to give the product. FAB ms: Calc.: 376 for C₂₂H₁₇N₂SOF; Obs.: 377 (M⁺+1).


EXAMPLES 45-49

- 15 The following compounds are prepared using the procedures described above. An alternative work up was utilized wherein the reaction mixture was concentrated in vacuo and the residue was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate and the combined organic phases were dried over MgSO₄, filtered and purified by flash chromatography eluting with 5% MeOH/CH₂Cl₂ to give
- 20 the product.


TABLE V			
Example #	Aryl ¹	Har	FAB ms
45	4-F-phenyl	4-(3-Me)-pyridyl	391
46	4-F-phenyl	4-quinolinyl	427
47	3-Cl-phenyl	4-pyridyl	393

- 61 -

48	4-F-phenyl	4-(2-Me)-pyridyl	391
49	3,4-F-phenyl	4-pyridyl	395

EXAMPLE 502-((4-methylsulfinylphenyl)-5-phenyl-3-(4-pyridyl) pyrrole

The title compound is prepared as in Example 44 using the product of Example 16 as the starting material. FAB ms: Calc.: 366 for C₂₃H₁₇N₂SO; Obs.: 367 (M⁺+1).

EXAMPLE 512-(phenyl)-5-(4-carboxyphenyl)-3-(4-pyridyl) pyrrole


To a solution of 25 mg (0.068 mmol) 2-phenyl-3-(4-pyridyl)-5-[(4-ethoxycarbonyl)phenyl]-1H-pyrrole (prepared according to Example 25) in 500 μ L methanol and 500 μ L THF was added 10 equivalents of a 5N NaOH solution. The reaction mixture was stirred at 60°C for 16 hours. After the mixture was cooled to room temperature, volatiles were removed *in vacuo*. The residue was taken up in methanol and acidified to pH 2 using a 2N HCl solution. Volatiles were removed *in vacuo*. The residue was dissolved in a 1:1 mixture of THF/methanol and tritrated. The resulting mixture was centrifuged and filtered to remove sodium chloride. This process was repeated 3 times to ensure total removal of sodium chloride. Solvents were evaporated *in vacuo* to afford the desired product as a white

- 62 -

glass, homogeneous by TLC (10% MeOH/CH₂Cl₂). FAB ms: Calc.: 340 for C₂₂H₁₆N₂O₂; Obs.: 341 (M⁺+1).

EXAMPLE 52

5




2-(phenyl)-5-(4-aminophenyl)-3-(4-pyridyl) pyrrole

To a suspension of 55 mg (0.161 mmol) 2-phenyl-3-(4-pyridyl)-5-[(4-nitro)phenyl]-1H-pyrrole (prepared according to Example 33) in 1.5 ml ethyl acetate and 1.5 ml ethanol was added 10 mg platinum(IV) oxide. The mixture was stirred under H₂ for 2 hours. The contents of flask were centrifuged, and the catalyst washed with ethyl acetate three times. The solvents were evaporated *in vacuo* to afford the desired product as a pale orange solid, homogeneous by TLC (5% MeOH/CH₂Cl₂). FAB ms: Calc.: 312 for C₂₁H₁₇N₃; Obs.: 313 (M⁺+1).

20


EXAMPLE 53



2-(phenyl)-5-(3-aminophenyl)-3-(4-pyridyl) pyrrole

The title compound is prepared as described above in Example 52 using the product of Example 27 as the starting material. FAB ms: Calc.: 312 for C₂₁H₁₇N₃; Obs.: 313 (M⁺+1).

- 63 -




EXAMPLE 542-(4-fluorophenyl)-5-(4-(2-pyridylmethylaminocarbonyl)-phenyl)-3-(4-pyridyl)-pyrrole

To a solution of 42 mg (0.124 mmol) 2-phenyl-3-(4-pyridyl)-5-[(4-carboxy)phenyl]-1H-pyrrole (prepared according to Example 51) in 800 μ L DMF was added 1.5 equivalents BOP reagent (benzotriazol-1-yloxy-tris-(dimethylamino)phosphonium hexafluorophosphate), 1.2 equivalents 2-(aminomethyl)pyridine, and 2.5 equivalents triethylamine. The mixture was stirred at room temperature for 15 hours. The reaction was quenched by the addition of 5% NaHCO₃ solution and ethyl acetate (EtOAc). After separation of phases, the aqueous phase was re-extracted with EtOAc. The combined organic layers were washed with 5% NaHCO₃ solution, water, brine, and dried over Na₂SO₄. After filtration and concentration of the filtrate *in vacuo*, the crude product was flash chromatographed (gradient elution with 1.0%-5.0% MeOH/CH₂Cl₂) to the desired product as a cream-colored solid, homogeneous by TLC. FAB ms: Calc.: 430 for C₂₈H₂₂N₄O; Obs.: 431 (M⁺+1).





EXAMPLES 55-63

This method is utilized to prepare the compounds listed below by coupling the appropriate amine to the acid derived from Example 51, or coupling an acid to the aniline formed in Example 52 or 53. The compounds are shown below in Table VI.



- 64 -

TABLE VI		
		
Example #	Aryl ² =	FAB ms
55	4-(CONHCH ₂ -phenyl)-phenyl 	430
56	4-(NHCO(CH ₂) ₂ -(1-piperidinyl))-phenyl 	451


- 65 -

57	<p>4-(CONHCH₂-4-pyr)-phenyl</p> 	431
58	<p>4-(CONH(CH₂)₂-(1-piperidinyl))-phenyl</p> 	451
59	<p>4-(CONH(CH₂)₂-2-(N-methylimidazolyl))phenyl</p> 	447
60	<p>4-(CONH(CH₂)₂-(2-pyridyl))-phenyl</p> 	445

- 66 -


61	<p data-bbox="634 247 1127 289">4-(CONHCH₂-(3-pyridyl))-phenyl</p> 	431
62	<p data-bbox="643 642 1110 684">3-(NHCO(CH₂)₃-NMe₂)-phenyl</p> 	425

- 67 -

EXAMPLE 635 2-(4-hydroxyphenyl)-5-(phenyl)-3-(4-pyridyl) pyrrole

13.2 mg of the product of Example 19 was stirred for 2 hours in a mixture of 50% trifluoroacetic acid in methylene chloride. The reaction mixture was concentrated in vacuo to provide the desired product. FAB ms: Calc.: 312 for C₂₁H₁₆N₂O; Obs.: 313 (M⁺+1).


10

EXAMPLE 6415 2-(3,4-dihydroxyphenyl)-5-(phenyl)-3-(4-pyridyl) pyrrole

20 mg of the product of Example 21 was dissolved in 1 mL of acetic acid and was hydrogenated at atmospheric pressure overnight in the presence of a catalytic amount of 10% Pd/C. The reaction mixture was concentrated in vacuo and the residue was purified by rotary


20 chromatography over silica gel eluting with a gradient of 5 to 10% MeOH/CH₂Cl₂ to provide the desired product. FAB ms: Calc.: 328 for C₂₁H₁₆N₂O₂; Obs.: 329 (M⁺+1).

- 68 -

EXAMPLE 652-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-3-(4-pyridyl) pyrrole

- 0.5 g (1.4 mmol) of the product of Example 44 was dissolved
5 in a mixture of 4 mL of methanol and 16 mL of ethyl acetate. The solution
was treated 0.045 g sodium tungstate dihydrate and 0.63 mL (5.6 mmol) of
30% hydrogen peroxide solution while heating at reflux over a period of 4
hours. A further 0.3 mL (2.8 mmol) hydrogen peroxide was added. The
mixture was refluxed overnight and then cooled to room temperature. A
10 white solid was recovered by filtration and was washed with water to
provide the desired product.

- $^1\text{H-NMR}$ (CDCl_3): 3.10 (s, 3H); 6.92 (d, 1H); 7.09 (t, 2H); 7.22 (m,
2H); 7.42 (dd, 2H); 7.72 (d, 2H); 7.95 (d, 2H); 8.45 (d, 2H), 8.95 (bs, 1H).
15 FAB ms: Calc: 392 for $\text{C}_{22}\text{H}_{17}\text{N}_2\text{SO}_2\text{F}$; Obs.: 393 (M^++1).

EXAMPLE 662-(3-chlorophenyl)-5-(4-methylsulfonylphenyl)-3-(4-pyridyl) pyrrole

20

The product of Example 47 was converted to the desired sulfone as described in Example 65. A portion of the product may be isolated by filtration as described in Example 65. The balance of the product was recovered following washing of the reaction mixture with aqueous


- 69 -

sodium sulfite, water and brine and drying over MgSO_4 . The crude product was purified by crystallization from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ followed by recrystallization from isopropanol.

- 5 $^1\text{H-NMR}$ (CD_3OD): 3.15 (s, 3H); 7.09 (d, 1H); 7.39 (m, 4H); 7.50 (s, 1H); 7.95 (s, 4H); 8.41 (d, 2H). FAB ms: Calc: 408 for $\text{C}_{22}\text{H}_{17}\text{N}_2\text{SO}_2\text{Cl}$; Obs.: 409 (M^++1).

EXAMPLES 67-123

- 10 Using the procedures set forth above, the following compounds can be prepared, as set forth in Table VII.


TABLE VII				
				
Ex#	$(\text{R}^a)_a\text{-Ar}^1$	$\text{Ar}^2\text{-(R}^a)_b$	$(\text{R}^a)_{0-3}\text{-HAr}$	R^2
67	Ph	Ph-4-F	3-quinolinyl	H
68	Ph	Ph-4-F	4-pyrimidinyl	H
69	Ph	Ph-4-F	3-pyridazinyl	H
70	Ph	Ph-4-F	2-pyrazinyl	H
71	Ph	Ph-4-CN	4-pyridyl	H
72	Ph	Ph-2-OMe	4-pyridyl	H
73	Ph	Ph-3-OMe	4-pyridyl	H
74	Ph-3,4-di-F	Ph-4-S(O)-Me	4-pyridyl	H
75	Ph	Ph-4-NMe ₂	4-pyridyl	H
76	Ph	4-(4-(N-COCH ₃) piperazinyl)-Ph	4-pyridyl	H
77	Ph	4-(morpholinyl)- Ph	4-pyridyl	H

- 70 -

78	Ph	Ph-2-Cl	4-pyridyl	H
79	Ph	Ph-3-Cl	4-pyridyl	H
80	Ph	Ph-4-CF ₃	4-pyridyl	H
81	Ph	Ph-4-S-Me	4-pyridyl	H
82	Ph	Ph-4-S(O)-Me	4-pyridyl	H
83	Ph-4-F	(4-methyl)-2-thiophenyl	4-pyridyl	H
84	Ph-4-F	(4-bromo)-2-thiophenyl	4-pyridyl	H
85	Ph-4-F	Ph-4-F-3-Cl	4-pyridyl	H
86	Ph-4-F	2-benzoxazolyl	4-pyridyl	H
87	Ph-4-F	2-benzofuranyl	4-pyridyl	H
88	Ph-4-F	4-(O(CH ₂) ₃ NMe ₂)-Ph	4-pyridyl	H
89	Ph-4-F	4-(O(CH ₂) ₂ -piperidin-1-yl)-Ph	4-pyridyl	H
90	Ph-4-Cl	Ph-4-F	4-pyridyl	H
91	Ph-3-Cl	Ph-4-F	4-pyridyl	H
92	Ph-2-Cl	Ph-4-F	4-pyridyl	H
93	Ph-3,4-di-Cl	Ph-4-F	4-pyridyl	H
94	Ph-3-CF ₃	Ph-4-F	4-pyridyl	H
95	Ph-4-S-Me	Ph-4-F	4-pyridyl	H
96	Ph-4-S(O)-Me	Ph-4-F	4-pyridyl	H
97	Ph-2-OBn	Ph-4-F	4-pyridyl	H
98	Ph-4-Br	Ph-4-F	4-pyridyl	H
99	Ph-2-OMe	Ph-4-F	4-pyridyl	H
100	Ph-3-OMe	Ph-4-F	4-pyridyl	H
101	Ph-4-OMe	Ph-4-F	4-pyridyl	H
102	Ph-4-NO ₂	Ph-4-F	4-pyridyl	H
103	Ph-4-NMe ₂	Ph-4-F	4-pyridyl	H
104	Ph-4-(4-N-COCH ₃)-piperazinyl	Ph-4-F	4-pyridyl	H
105	Ph-4-morpholinyl	Ph-4-F	4-pyridyl	H
106	2-thiophenyl	Ph-4-F	4-pyridyl	H
107	3-thiophenyl	Ph-4-F	4-pyridyl	H
108	2-furoyl	Ph-4-F	4-pyridyl	H
109	3-furoyl	Ph-4-F	4-pyridyl	H
110	Ph-3-Cl	Ph-4-S(O)Me	4-(2-Me)-pyridyl	H

- 71 -

111	Ph-3,4-di-F	Ph-4-S(O)Me	4-(2-Me)-pyridyl	H
112	Ph-3,4-di-F	Ph-4-S(O)Me	4-pyridyl	F
113	Ph-3-CF ₃	Ph	4-pyridyl	H
114	Ph-4-F	Ph-4-S(O)Me	4-(2-amino benzyl)pyridyl	H
115	3-quinolinyl	Ph	4-pyridyl	H
116	Ph-4-F	4-(SMe)-Ph	4-pyridyl	Br
117	Ph-3,4-di-F	Ph	4-pyridyl	CO ₂ Et
118	Ph-3,4-di-F	Ph	4-pyridyl	CN
119	3-(CF ₃)-Ph	Ph	4-pyridyl	CO ₂ Et
120	3-(CF ₃)-Ph	Ph	4-pyridyl	CN
121	Ph	Ph-4-F	3-methyl-4- pyridyl	H
122	Ph-3-Cl	Ph-4-S(O)Me	4-(2-amino) pyrimidinyl	H
123	Ph	Ph-4-F	3,5-dimethyl-4- pyridyl	H
Bn = benzyl, Ph = phenyl, Me = methyl Et = ethyl .				


EXAMPLE 124

5

A mixture of ethyl 2-benzoyl-acetate (0.41 g (2.0 mmol), 0.5 g (1.44 mmol) of the product of Preparative Example 61 and 0.61 g (8 mmol) of ammonium acetate were heated in acetic acid at reflux until the benzoin was consumed. The reaction mixture was diluted with ethyl acetate and washed with water and brine and dried over MgSO₄. The mixture was filtered and the filtrate was concentrated in vacuo and the residue was purified by chromatography over silica gel to give the desired product. ¹H-NMR (CDCl₃, 300 MHz): 0.92 (t, 3H); 4.01 (q, 2H); 6.94 (t, 2H); 7.12 (m, 4H); 7.41 (m, 4H); 7.61 (m, 2H); 8.10 (m, 1H); 9.20 (bs, 1H). FAB ms: C₂₄H₁₉N₂O₂F=386; Observed: 387 (M⁺=1).

15


- 72 -

EXAMPLE 125

5

A mixture of benzyl 2-benzoyl-acetate (2.0 mmol), 0.5 g (1.44 mmol) of the product of Preparative Example 61 and 0.61 g (8 mmol) of ammonium acetate heated in acetic acid at reflux will provide the title compound after purification as recited in Example 124.

10


EXAMPLE 126

15


A mixture of benzyl 2-(4-fluorophenyl)-3-(4-pyridyl)-5-phenylpyrrole-4-carboxylate from Example 125 (1.0 mmol), 0.01 g of 10% Pd/C in 5 mL of EtOH will yield 2-(4-fluorophenyl)-3-(4-pyridyl)-5-phenylpyrrole-4-carboxylic acid after treatment with 40 psi H₂ followed by filtration.

20

- 73 -


EXAMPLE 127

- 5 A mixture of 2-(4-fluorophenyl)-3-(4-pyridyl)-5-phenyl-
pyrrole-4-carboxylic acid from Example 126 (1.0 mmol), EDC, Hunig's
base and dimethylamine hydrochloride in DMF is stirred at room
temperature overnight. The reaction mixture is diluted with water, adjusted
to pH 7.0 and extracted with ethyl acetate. The organic extracts are washed
10 with brine, dried over MgSO₄, filtered and concentrated in vacuo. The
residue is purified by chromatography.

EXAMPLE 128

- 15 A mixture of benzoylacetone 0.5 g (3.4 mmol), 1.17 g (3.4
mmol) of the product of Preparative Example 61 and 1.0 g (13.6 mmol)
ammonium acetate are heated in acetic acid at reflux until the benzoin is
20 consumed. The reaction mixture was diluted with ethyl acetate and washed
with water and brine and dried over MgSO₄. The mixture is filtered and the
filtrate is concentrated in vacuo. The residue is purified by chromatography
over silica gel eluting with 5% MeOH/CH₂Cl₂ to give the desired product.
H¹-NMR (CDCl₃, 300 MHz): 7.0 (t, 2H); 7.24 (m, 2H); 7.32-7.48 (m, 5H);
25 7.74 (bd, 2H); 8.42 (bs, 1H). FAB ms: C₂₂H₁₄N₃F = 339; Observed: 340
(M⁺=1).

- 74 -

EXAMPLE 129


5

The product of Example 5 is dissolved in methylene chloride and treated with 1.05 equivalents of n-propylsulfinyl chloride at 0°C under nitrogen. After 30 minutes triethylamine is added to neutralize the reaction mixture. The reaction mixture is diluted with ethyl acetate and washed with water and brine and dried over MgSO₄. The mixture is filtered and the filtrate is concentrated in vacuo. The residue is purified by chromatography over silica gel to give the desired product.

10

EXAMPLE 130


15



The product of Example 129 is reacted with 1.05 equivalents of meta-chloroperoxybenzoic acid in CH₂Cl₂ at 0°C. The reaction mixture is stirred overnight at room temperature. The solution is diluted with EtOAc and washed with saturated sodium bicarbonate solution followed by brine. The solution is dried over MgSO₄, filtered and concentrated in vacuo. The residue is purified by silica gel chromatography to produce the desired product.

20
25EXAMPLE 131


- 75 -




To 5 ml of DMF at room temperature under nitrogen is added 0.3 g (2 mmol) of POCl₃ dropwise. After 15 minutes a solution of 0.37 g (0.86 mmol) of the product of Example 15 is added dropwise. The solution was warmed at 60°C until the starting material had been consumed. The reaction mixture was cooled to room temperature and then poured into ice water (20 ml). The mixture was made basic by addition of saturated sodium carbonate solution and then stirred in the presence of 20 ml of chloroform. The chloroform phase was separated and the aqueous phase was extracted with chloroform (2 x 10 ml). The combined organic phase is washed with water and brine and dried over MgSO₄. The mixture is filtered and the filtrate is concentrated in vacuo; the residue is purified by chromatography over silica gel to give the desired product.

15

- 76 -

EXAMPLE 132

5 The product of Example 131 is dissolved in t-butyl alcohol and methyl-2-butene (6:1 ratio). The solution is then treated with 1.5 eq of monobasic sodium phosphate and an aqueous solution of sodium chlorate. The reaction mixture is stirred at room temperature until the starting
10 product is extracted with ethyl acetate and the combined organic phase is washed with water and brine, and dried over MgSO₄. The mixture is filtered and the filtrate is concentrated in vacuo to give the desired product.

EXAMPLE 133

15 To 5 ml of N,N-dimethylbutyramide at room temperature under nitrogen is added 0.3 g (2 mmol) of POCl₃ dropwise. After 15 minutes, a solution of 0.37 g (0.86 mmol) of the product of Example 4 is
20 added dropwise. The solution is warmed at 60°C until the starting material is consumed. The reaction mixture is cooled to room temperature and then poured into ice water (20 ml). The mixture is made basic by addition of saturated sodium carbonate solution and then stirred in the presence of 20 mL of chloroform. The chloroform phase is separated and the aqueous
25 phase extracted with chloroform (2 x 10 ml). The combined organic phase


- 77 -

is washed with water and brine and dried over MgSO_4 . The mixture is filtered and the filtrate is concentrated in vacuo. The residue is purified by chromatography over silica gel to give the desired product.

5

EXAMPLES 134-203

Using the procedures set forth above, the compounds shown in Table VIII can be prepared.

TABLE VIII				
				
Ex.	$(R^a)_a$ -Ar ¹	Ar ² -(R ^a) _b	$(R^a)_{0-3}$ -HAr	R ²
134	Ph-4-F	Ph-4-OMe	4-pyridyl	C(O)Me
135	Ph-4-F	Ph-2,5-di-OMe	4-pyridyl	C(O)Me
136	Ph-4-F	Ph-4-Br	4-pyridyl	C(O)Propyl
137	Ph-4-F	Ph-4-Cl	4-pyridyl	C(O)Ethyl
138	Ph-4-F	Ph-4-OMe	2-pyridyl	C(O)Me
139	Ph-4-F	Ph-4-Br	2-pyridyl	C(O)Me
140	Ph-4-F	Ph-2,5-di-OMe	2-pyridyl	C(O)Me
141	Ph	Ph-4-OMe	4-pyridyl	C(O)Me
142	Ph	Ph-4-Cl	4-pyridyl	C(O)Me
143	Ph	Ph-2,5-di-OMe	4-pyridyl	C(O)Me
144	Ph	Ph-4-F	4-pyridyl	C(O)Me
145	Ph-4-Cl	Ph	4-pyridyl	C(O)Me
146	Ph-4-F	Ph-4-SMe	4-pyridyl	C(O)Me
147	Ph-4-SMe	Ph	3-methyl-4-pyridyl	C(O)Me

- 78 -

148	Ph-4-F	Ph-4-SMe	3-methyl-4-pyridyl	CN
149	Ph-4-F	Ph-4-S(O)Me	3-methyl-4-pyridyl	CN
150	Ph-3-Cl	Ph-4-Cl	4-quinolyl	CN
151	Ph-4-F	Ph-4-Cl	2-methyl-4-pyridyl	CN
152	Ph	Ph-4-F	3,5-dimethyl-4-pyridyl	CN
153	Ph	Ph-4-F	3-quinolinylnyl	CN
154	Ph	Ph-4-F	4-quinolinylnyl	CN
155	Ph	Ph-4-F	2-quinolinylnyl	CN
156	Ph	Ph-4-F	2-pyrimidinyl	CN
157	Ph	Ph-4-F	4-pyrimidinyl	CN
158	Ph	Ph-4-F	3-pyridazinyl	CN
159	Ph	Ph-4-F	2-pyrazinyl	CN
160	Ph	Ph-4-F	2-pyrimidinyl	CN
161	Ph	Ph-4-F	4-pyrimidinyl	CN
162	Ph	Ph-4-F	2-imidazo-(4,5-b)-pyridinyl	CN
163	Ph	Ph-4-F	7-imidazo-(4,3-b)-pyridinyl	CN
164	Ph	Ph-4-F	4-pyridyl	COMe
165	Ph	Ph-4-F	4-pyridyl	SO ₂ Me
166	Ph	Ph-4-CN	4-pyridyl	COMe
167	Ph	Ph-2-OMe	4-pyridyl	COMe
168	Ph	Ph-3-OMe	4-pyridyl	CN
169	Ph	Ph-4-OMe	4-pyridyl	CO ₂ Et
170	Ph	Ph-4-NO ₂	4-pyridyl	CN
171	Ph	Ph-4-NMe ₂	4-pyridyl	CN
172	Ph	4-(4-(N-COCH ₃)-piperazinyl)-Ph	4-pyridyl	CN
173	Ph	4-(morpholinyl)-Ph	4-pyridyl	CN
174	Ph	Ph-2-Cl	4-pyridyl	CN

- 79 -

175	Ph	Ph-3-Cl	4-pyridyl	C(O)Me
176	Ph	Ph-4-CF ₃	4-pyridyl	SO ₂ Me
177	Ph	Ph-4-S-Me	4-pyridyl	CN
178	Ph	Ph-4-S(O)-Me	4-pyridyl	CN
179	Ph-4-F	(4-methyl) thiophen-2yl	4-pyridyl	CN
180	Ph-4-F	(3-methyl) thiophen-2yl	4-pyridyl	C(O)Me
181	Ph-4-F	(4-bromo) thiophen-2yl	4-pyridyl	SO ₂ Me
182	Ph-4-F	(5-methyl) thiophen-2yl	4-pyridyl	CN
183	Ph-4-F	Ph-4-F-3-Cl	4-pyridyl	CN
184	Ph-4-F	2-benzoxazolyl	4-pyridyl	CN
185	Ph-4-F	2-benzofuranyl	4-pyridyl	CN
186	Ph-4-F	4-(O(CH ₂) ₃ - NMe ₂)-Ph	4-pyridyl	CN
187	Ph-4-F	4-(O(CH ₂) ₂ N- piperidiny)-Ph	4-pyridyl	CN
188	Ph-4-Cl	Ph-4-F	4-pyridyl	CN
189	Ph-3-Cl	Ph-4-F	4-pyridyl	CN
190	Ph-2-Cl	Ph-4-F	4-pyridyl	CN
191	Ph-3,4-di-Cl	Ph-4-F	4-pyridyl	CN
192	Ph-3-CF ₃	Ph-4-F	4-pyridyl	CN
193	Ph-4-S-Me	Ph-4-F	4-pyridyl	CN
194	Ph-4-S(O)-Me	Ph-4-F	4-pyridyl	CN
195	Ph-2-OBn	Ph-4-F	4-pyridyl	CN
196	Ph-4-Br	Ph-4-F	4-pyridyl	CN
197	Ph-2-OMe	Ph-4-F	4-pyridyl	CN
198	Ph-3-OMe	Ph-4-F	4-pyridyl	CN
199	Ph-4-OMe	Ph-4-F	4-pyridyl	CN
200	Ph-4-NO ₂	Ph-4-F	4-pyridyl	CN
201	Ph-4-NMe ₂	Ph-4-F	4-pyridyl	CN

- 80 -

202	4-(4-(N-COCH ₃)-piperazinyl)-Ph	Ph-4-F	4-pyridyl	CN
203	4-(morpholinyl)-Ph	Ph-4-F	4-pyridyl	CN
Me = methyl, Et = ethyl, Ph = phenyl, Bn = benzyl,				

BIOLOGICAL ASSAYS.

The ability of compounds of the present invention to inhibit cytokines can be demonstrated by the following *in vitro* assays.

5

Lipopolysaccharide mediated production of cytokines

Human peripheral blood mononuclear cells (PBMC) are isolated from fresh human blood according to the procedure of Chin and Kostura, *J. Immunol.* **151**, 5574-5585 (1993). Whole blood is collected by sterile venipuncture into 60 mL syringes coated with 1.0 mL of sodium-heparin (Upjohn, 1000 U/mL) and diluted 1:1 in Hanks Balanced Salt Solution (Gibco). The erythrocytes are separated from the PBMC's by centrifugation on a Ficoll-Hypaque lymphocyte separation media. The PBMC's are washed three times in Hanks Balanced Salt Solution and then resuspended to a final concentration of 2×10^6 cell/mL in RPMI (cell culture medium) containing 10% fresh autologous human serum, penicillin streptomycin (10 U/mL) and 0.05% DMSO. Lipopolysaccharide (Salmonella type Re545; Sigma Chemicals) is added to the cells to a final concentration of 100 ng/mL. An aliquot (0.1 mL) of the cells is quickly dispensed into each well of a 96 well plate containing 0.1 mL of the test compound, at the appropriate dilution, and incubated for 24 hours. at 37°C in 5% CO₂. At the end of the culture period, cell culture supernatants are assayed for IL-1 β , TNF- α , IL-6 and PGE₂ production using specific ELISA.

25

IL-1 mediated cytokine production

Human peripheral blood mononuclear cells are isolated from fresh human blood according to the procedure of Chin and Kostura, *J. Immunol.* **151**, 5574-5585 (1993). Whole blood is collected by sterile venipuncture into 60 mL syringes coated with 1.0 mL of sodium-heparin (Upjohn, 1000 U/mL) and diluted 1:1 in Hanks Balanced Salt Solution (Gibco). The erythrocytes are separated from the PBMC's by

30

- 81 -

centrifugation on a Ficoll-Hypaque lymphocyte separation media. The PBMC's are washed three times in Hanks Balanced Salt Solution and then resuspended to a final concentration of 2×10^6 cell/mL in RPMI containing 10% fresh autologous human serum, penicillin streptomycin (10 U/mL) and 0.05% DMSO. Endotoxin free recombinant human IL-1 β is then added to a final concentration of 50 pMolar. An aliquot (0.1 mL) of the cells is quickly dispensed into each well of a 96 well plate containing 0.1 mL of the compound at the appropriate dilution. and are incubated for 24 hours at 37°C in 5% CO₂. At the end of the culture period, cell culture supernatants are assayed for TNF- α , IL-6 and PGE₂ synthesis using specific ELISA.

Determination of IL-1 β , TNF- α , IL-6 and prostanoid production from LPS or IL-1 stimulated PBMC's

IL-1 β ELISA.

Human IL-1 β can be detected in cell-culture supernatants or whole blood with the following specific trapping ELISA. Ninety-six well plastic plates (Immulon 4; Dynatech) are coated for 12 hours at 4°C with 1 mg/mL protein-A affinity chromatography purified mouse anti-human IL-1b monoclonal antibody (purchased as an ascites preparation from LAO Enterprise, Gaithersburg Maryland) diluted in Dulbecco's phosphate-buffered saline (-MgCl₂, -CaCl₂). The plates are washed with PBS (phosphate buffered saline)-Tween (Kirkegaard and Perry) then blocked with 1% BSA diluent and blocking solution (Kirkegaard and Perry) for 60 minutes at room temperature followed by washing with PBS Tween.

IL-1 β standards are prepared from purified recombinant IL-1 β produced from *E. coli*. The highest concentration begins at 10 ng/mL followed by 11 two-fold serial dilutions. For detection of IL-1 β from cell culture supernatants or blood plasma, 10 - 25 mL of supernatant is added to each test well with 75 - 90 mL of PBS Tween. Samples are incubated at room temperature for 2 hours then washed 6 times with PBS Tween on an automated plate washer (Dennly). Rabbit anti-human IL-1 β polyclonal antisera diluted 1:500 in PBS-Tween is added to the plate and incubated for 1 hour at room temperature followed by six washes with PBS-Tween. Detection of bound rabbit anti-IL-1 β IgG is accomplished with Fab' fragments of Goat anti-rabbit IgG-horseradish peroxidase conjugate

- 82 -

(Accurate Scientific) diluted 1:10,000 in PBS-Tween. Peroxidase activity was determined using TMB peroxidase substrate kit (Kirkegaard and Perry) with quantitation of color intensity on a 96-well plate Molecular Devices spectrophotometer set to determine absorbance at 450 nM. Samples are
5 evaluated using a standard curve of absorbance versus concentration. Four-parameter logistics analysis generally is used to fit data and obtain concentrations of unknown compounds.

TNF- α ELISA

10 Immulon 4 (Dynatech) 96-well plastic plates are coated with a 0.5 mg/mL solution of mouse anti-human TNF- α monoclonal antibody. The secondary antibody is a 1:2500 dilution of a rabbit anti-human TNF- α polyclonal serum purchased from Genzyme. All other operations are identical to those described above for IL-1 β . The standards are prepared in
15 PBS-Tween + 10% FBS (fetal bovine serum) or HS (human serum). Eleven 2 fold dilutions are made beginning at 20 ng/mL TNF- α .

IL-6 ELISA

20 Levels of secreted human IL-6 are also determined by specific trapping ELISA as described previously in Chin and Kostura, *J. Immunol.* **151**, 5574-5585 (1993). (Dynatech) ELISA plates are coated with mouse anti-human IL-6 monoclonal antibody diluted to 0.5 mg/ml in PBS. The secondary antibody, a rabbit anti-human IL-6 polyclonal antiserum, is diluted 1:5000 with PBS-Tween. All other operations are identical to those
25 described above for IL-1 β . The standards are prepared in PBS-Tween + 10% FBS or HS. Eleven 2 fold dilutions are made beginning at 50 ng/mL IL-6.

PGE₂ production

30 Prostaglandin E₂ is detected in cell culture supernatants from LPS or IL-1 stimulated PBMC's using a commercially available enzyme immunoassay. The assay purchased from the Cayman Chemical (Catalogue number 514010) and is run exactly according to the
35 manufacturers instructions.


Interleukin8 (IL-8)

- 83 -

The present compounds can also be assayed for IL-8 inhibitory activity as discussed below. Primary human umbilical cord endothelial cells (HUVEC) (Cell Systems, Kirland, Wa.) are maintained in culture medium supplemented with 15% fetal bovine serum (FBS) and 1% CS-HBGF (cell culture additive) consisting of aFGF (acid fibroblast growth factor) and heparin. The cells are then diluted 20-fold before being plated (250 μ l) into gelatin coated 96-well plates. Prior to use, culture medium is replaced with fresh medium (200 μ l). Buffer or test compound (25 μ l, at appropriate concentrations) is then added to each well in quadruplicate wells and the plates incubated for 6h in a humidified incubator at 37°C in an atmosphere of 5% CO₂. At the end of the incubation period, supernatant is removed and assayed for IL-8 concentration using an IL-8 ELISA kit obtained from R&D Systems (Minneapolis, MN). All data is presented as mean value (ng/ml) of multiple samples based on the standard curve. IC₅₀ values where appropriate are generated by non-linear regression analysis.

The following compounds are found to inhibit cytokines at IC₅₀ concentrations of less than 100 μ M.

TABLE IX



R^1	$(R^a)_a - Ar^1$	$Ar^2 - (R^a)_b$	$(R^a)_{0-3} - HAr$	R^2
H	4-F-Ph	4-F-Ph	4-Pyr	H
H	4-F-Ph	4-OMe-Ph	4-Pyr	H
H	4-F-Ph	2,5-di-(OMe)-Ph	4-Pyr	H
H	4-F-Ph	4-Br-Ph	4-Pyr	H
H	4-F-Ph	4-Cl-Ph	4-Pyr	H
H	Ph	4-OMe-Ph	4-Pyr	H
H	Ph	4-Cl-Ph	4-Pyr	H
H	Ph	2,5-di-(OMe)-Ph	4-Pyr	H
H	Ph	4-F-Ph	4-Pyr	H
H	4-F-Ph	4-SMe-Ph	4-Pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-Pyr	H
H	4-Cl-Ph	Ph	4-Pyr	H
H	4-SMe-Ph	Ph	4-Pyr	H
H	4-S(O)Me-Ph-4-Pyr	Ph	H	Ph
H	4-CF ₃ -Ph	Ph	4-Pyr	H
H	4-Me-Ph	Ph	4-Pyr	H
H	4-OH-Ph	Ph	4-Pyr	H
H	3,4-di-Cl-Ph	Ph	4-Pyr	H
H	3,4-di-OH-Ph	Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	CO ₂ Et

- 85 -

H	3-Cl-Ph	Ph	4-pyr	H
H	4-CN-Ph	Ph	4-Pyr	H
H	Ph	4-CO ₂ Et-Ph	4-Pyr	H
H	Ph	2-F-Ph	4-Pyr	H
H	Ph	3-NO ₂ -Ph	4-Pyr	H
H	5-Me-thiophen-2-yl	Ph	4-Pyr	H
H	3-Me-thiophen-2-yl	Ph	4-Pyr	H
H	3-quinolinyl	Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	CN
H	4-F-Ph	4-(SMe)-Ph	4-(3-Me)-Pyr	H
H	4-F-Ph	4-(SMe)-Ph	4-(3-Me)-Pyr	H
H	Ph	2-(NO ₂)-Ph	4-Pyr	H
H	Ph	3-F-Ph	4-Pyr	H
H	Ph	4-CO ₂ H-Ph	4-Pyr	H
H	Ph	3-NH ₂ -Ph	4-Pyr	H
H	Ph	4-NO ₂ -Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	H
H	2,4-di-F-Ph	Ph	4-Pyr	H
H	3-CN-Ph	Ph	4-Pyr	H
H	3,4-di-F-Ph	Ph	4-Pyr	H
H	Ph	4-NH ₂ -Ph	4-Pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-quinolinyl	H
H	4-F-Ph	4-SMe-Ph	4-pyr	Br
H	4-F-Ph	4-SMe-Ph	4-quinolinyl	H
H	Ph	4-(CONHCH ₂ -phenyl)-Ph	4-pyr	H
H	Ph	4-(NHCO(CH ₂) ₃ -NMe ₂)-Ph	4-pyr	H
H	Ph	4-(CONHCH ₂ -4-pyr)-Ph	4-pyr	H
H	Ph	4-(CONH(CH ₂) ₂ -(1-piperidinyl))-Ph	4-pyr	H
H	Ph	4-(CONH(CH ₂) ₂ -2-(N-methylimidazolyl))-Ph	4-pyr	H
H	Ph	4-(CONHCH ₂ -(2-pyridyl))-Ph	4-pyr	H
H	Ph	4-(CONH(CH ₂) ₂ -(2-pyridyl))-Ph	4-pyr	H


- 86 -

H	Ph	4-(CONHCH ₂ -(3-pyridyl))-Ph	4-pyr	H
H	Ph	3-(NHCO(CH ₂) ₃ NMe ₂)-Ph	4-pyr	H
H	Ph	4-(NHCO(CH ₂) ₂ -(1-piperidyl))-Ph	4-pyr	H
H	2-Cl-Ph	Ph	4-pyr	H
H	3-Cl-Ph	4-SMe-Ph	4-pyr	H
H	4-F-Ph	Ph	4-pyrimidinyl	F
H	4-F-Ph	4-SMe-Ph	4-(2-methyl)-pyr	H
H	3-Cl-Ph	4-S(O)Me-Ph	4-pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-(2-methyl)-pyr	H
H	3,4-di-F-Ph	4-SMe-Ph	4-pyr	H
H	3,4-di-F-Ph	4-S(O)Me-Ph	4-Pyr	H
H	4-F-Ph	4-SO ₂ Me-Ph	4-pyr	H
H	3,4-di-F-Ph	Ph	4-pyr	CO ₂ Et
H	3,4-di-F-Ph	Ph	4-pyr	CN
H	3-CF ₃ -Ph	Ph	4-pyr	CO ₂ Et
H	3-CF ₃ -Ph	Ph	4-pyr	CN
H	3-Cl-Ph	4-SO ₂ Me-Ph	4-pyr	H


- 87 -

WHAT IS CLAIMED IS:

1. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:



and each independently represent a 5-10 membered aryl or heteroaryl group substituted with R^a groups;

wherein a and b represents integers, 0, 1, 2 or 3, such that the sum of a plus b is 1, 2, 3 or 4;



represents a heteroaryl group containing from 5 to 10 atoms, 1-4 of which are heteroatoms, 0-4 of which heteroatoms are N and 0-1 of which are O or S, said heteroaryl group being unsubstituted or substituted with 0-3 R^a groups; each R^a independently represents a member selected from the group consisting of: halo; CN, NO₂, R²¹; OR²³; SR²³; S(O)R²¹; SO₂R²¹; NR²⁰R²³; NR²⁰COR²¹; NR²⁰CO₂R²¹; NR²⁰CONR²⁰R²³; NR²⁰SO₂R²¹; NR²⁰C(NR²⁰)NHR²⁰, CO₂R²³; CONR²⁰R²³; SO₂NR²⁰R²³; SO₂NR²⁰COR²¹; SO₂NR²⁰CONR²⁰R²³; SO₂NR²⁰CO₂R²¹; OCONR²⁰R²³; OCONR²⁰SO₂R²⁰, C(O)OCH₂OC(O)R²⁰; C(NR²⁰)NR²⁰R²³ and CONR²⁰SO₂R²¹;

R¹ is selected from the group consisting of: H, aryl, C₁₋₁₅ alkyl, C₃₋₁₅ alkenyl, C₃₋₁₅ alkynyl and heterocyclyl, said alkyl, aryl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one

- 88 -

to three members selected from the group consisting of: aryl, heteroaryl, heterocyclyl, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{21}$, SO_2R^{21} , $SO_2NR^{20}R^{23}$, $SO_2NR^{20}COR^{21}$, $SO_2NR^{20}CONR^{20}R^{23}$, $NR^{20}COR^{21}$, $NR^{20}CO_2R^{21}$, $NR^{20}CONR^{20}R^{23}$, $N(R^{20})C(NR^{20})NHR^{20}$, CO_2R^{20} , $CONR^{20}R^{23}$,
 5 $CONR^{20}SO_2R^{21}$, $NR^{20}SO_2R^{21}$, $SO_2NR^{20}CO_2R^{21}$, $CONR^{20}R^{23}$, $CONR^{20}SO_2R^{21}$; $CONR^{20}R^{23}$ and $C(O)OCH_2OC(O)R^{20}$;

R^2 is selected from the group consisting of: H, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, halo, NO_2 , heterocyclyl, CN, $S(O)R^{21}$,
 10 SO_2R^{21} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{21}$, $SO_2NR^{20}CON(R^{20})_2$, COR^{20} , CO_2R^{20} , $CONR^{20}R^{23}$, $CONR^{20}SO_2R^{21}$ and $SO_2NR^{20}CO_2R^{21}$, said alkyl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: halo, heterocyclyl, CN, aryl, heteroaryl, R^{20} , OR^{20} , SR^{20} , $NR^{20}R^{23}$, $S(O)R^{21}$, SO_2R^{21} ,
 15 $SO_2NR^{20}R^{23}$, $SO_2NR^{20}COR^{21}$, $SO_2NR^{20}CONR^{20}R^{23}$, $NR^{20}COR^{21}$, $NR^{20}CO_2R^{21}$, $NR^{20}CONR^{20}R^{23}$, $NR^{20}C(NR^{20})NHR^{20}$, CO_2R^{20} , $CONR^{20}R^{23}$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{21}$, $SO_2NR^{20}CO_2R^{22}$, $CONR^{20}SO_2R^{21}$ and $CONR^{20}R^{23}$;

20 R^{20} represents a member selected from the group consisting of: H, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl, alkenyl, alkynyl, heterocyclyl, aryl and heteroaryl being optionally substituted with 1-3 groups selected from halo, aryl and heteroaryl;

25 R^{21} represents a member selected from the group consisting of: C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl, such alkyl, alkenyl and alkynyl being optionally interrupted with oxo and/or 1-2 heteroatoms selected from O, S, $S(O)$, SO_2 and NR^{20} , said alkyl, alkenyl, alkynyl, aryl and heteroaryl being optionally substituted with from 1-3 of halo, heterocyclyl, aryl, heteroaryl, CN, OR^{20} , $O((CH_2)_nO)_mR^{20}$, $NR^{20}((CH_2)_nO)_mR^{20}$ wherein n represents an integer of from 2 to 4, and m represents an integer of from 1 to 3; SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $NR^{20}COR^{22}$,
 30 $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{22}C(NR^{22})NHR^{22}$, CO_2R^{20} ,

- 89 -

CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²²,
OCONR²⁰SO₂R²²OC(O)R²⁰, C(O)OCH₂OC(O)R²⁰ and OCON(R²⁰)₂;

5 R²² is selected from the group consisting of: C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl, alkenyl, and alkynyl being optionally substituted with 1-3 halo, aryl or heteroaryl groups;

10 R²³ is R²¹ or H;

R²⁴ is selected from COR²², CO₂R²², CON(R²⁰)₂, SO₂R²² and R²³;

and when two R²⁰ groups are present, when R²⁰ and R²¹ are present, or when R²⁰ and R²³ are present, said two R²⁰ groups, R²⁰ and R²¹ or said R²⁰ and R²³ may be taken in combination with the atoms to which they are attached and any intervening atoms and represent heterocyclyl containing from 5-10 atoms, at least one atom of which is a heteroatom selected from O, S or N, said heterocyclyl optionally containing 1-3 additional N atoms and 0-1 additional O or S atom.

20

2. A compound in accordance with claim 1 wherein:
Ar¹ and Ar² are independently and selected from the group consisting of:

- a) phenyl,
- b) pyridyl,
- 25 c) pyrimidinyl,
- d) thiophenyl,
- e) furanyl,
- f) imidazolyl,
- g) thiazolyl,
- 30 h) isothiazolyl,
- i) oxazolyl,
- j) isoxazolyl, and
- k) naphthyl.

35

3. A compound in accordance with claim 1 wherein
HAr is selected from the group of consisting of:

- 90 -

- 5
- a) pyridyl,
 - b) quinolyl,
 - c) purinyl,
 - d) imidazolyl,
 - e) imidazopyridyl,
 - f) pyrimidinyl and
 - g) pyridazinyl.

10 4. A compound in accordance with claim 1 wherein R¹ is hydrogen.

5 A compound in accordance with claim 1 wherein R¹ is substituted or unsubstituted C₁₋₁₅ alkyl.

15 6. A compound in accordance with claim 1 wherein R² represents a member selected from the group consisting of:

- 20
- a) H;
 - b) C₁₋₆ alkyl;
 - c) halo;
 - d) CN;
 - e) C(O)C₁₋₆ alkyl;
 - f) C(O)C₁₋₆ alkylphenyl;
 - g) CO₂H;
 - h) CO₂C₁₋₆ alkyl
 - 25 i) CO₂C₁₋₆ alkylphenyl;
 - j) CONH₂;
 - k) CONHC₁₋₆ alkyl;
 - l) C(O)N(C₁₋₆ alkyl)₂;
 - m) SO₂NH₂;
 - 30 n) SO₂NHC₁₋₆ alkyl and
 - o) SO₂N(C₁₋₆ alkyl)₂.

35 7. A compound in accordance with claim 1 wherein: Ar¹ and Ar² are independently selected from:

- a) phenyl,
- b) pyridyl,

- 91 -

- 5 c) pyrimidinyl,
 d) thiophenyl,
 e) furanyl,
 f) imidazolyl,
 g) thiazolyl,
 h) isothiazolyl,
 i) oxazolyl,
 j) isoxazolyl and
 k) naphthyl;
- 10 one, two or three R^a groups are present and attached to Ar^1 and Ar^2 , and each R^a is independently selected from the group consisting of: halo, R^{21} , OR^{23} , $NR^{20}R^{23}$, CO_2R^{23} , $CONR^{20}R^{23}$, SO_2R^{21} and $S(O)R^{21}$, R^{20} , R^{21} and R^{23} are as originally defined;
- 15 HAr is selected from:
- a) pyridyl,
 b) quinolyl,
 c) purinyl,
 d) imidazolyl,
 20 e) imidazopyridyl,
 f) pyrimidinyl and
 g) pyridazinyl;
- R^1 is: a) H or
 25 b) substituted or unsubstituted alkyl; and
- R^2 is selected from the group consisting of:
- a) H,
 b) alkyl,
 30 c) halo,
 d) CN,
 e) $C(O)C_{1-6}$ alkyl,
 f) $C(O)C_{1-6}$ alkylphenyl,
 g) CO_2H ,
 35 h) CO_2C_{1-6} alkyl,
 i) CO_2C_{1-6} alkylphenyl,

- 92 -

- 5
- j) CONH_2 ,
 - k) CONHC_{1-6} alkyl,
 - l) $\text{C(O)N(C}_{1-6} \text{ alkyl)}_2$,
 - m) SO_2NH_2 ,
 - n) $\text{SO}_2\text{NHC}_{1-6}$ alkyl and
 - o) $\text{SO}_2\text{N(C}_{1-6} \text{ alkyl)}_2$.

8. A compound in accordance with claim 1 wherein:
 $(\text{Ra})_a\text{-Ar}^1$ is selected from the group consisting of:

- 10
- a) phenyl,
 - b) 4-fluorophenyl,
 - c) 4-chlorophenyl,
 - d) 3-fluorophenyl,
 - e) 3-chlorophenyl,
 - 15 f) thiophen-2-yl,
 - g) thiophen-3-yl,
 - h) 4-fluorothiophen-2-yl,
 - i) 4-fluorothiophen-3-yl,
 - j) 5-fluorothiophen-2-yl,
 - 20 k) 5-fluorothiophen-3-yl,
 - l) 4-chlorothiophen-2-yl,
 - m) 4-chlorothiophen-3-yl,
 - n) 5-chlorothiophen-2-yl,
 - o) 5-chlorothiophen-3-yl,
 - 25 p) 3-methylphenyl,
 - q) 3,4-dichlorophenyl,
 - r) 3-hydroxyphenyl,
 - s) 4-hydroxyphenyl,
 - t) 3,4-dihydroxyphenyl,
 - 30 u) 3-methyl-2-thiophenyl,
 - v) 5-methyl-2-thiophenyl,
 - w) 4-carboxymethylphenyl,
 - x) 3-cyanophenyl,
 - y) 4-cyanophenyl,
 - 35 z) 2-pyridyl,
 - aa) 2-furoyl,

- 93 -

- 5
10
15
20
- bb) 3-furoyl,
 - cc) 4-methylsulfinylphenyl,
 - dd) 4-trifluoromethylphenyl,
 - ee) 3-trifluoromethylphenyl,
 - ff) 4-methylphenyl,
 - gg) 4-t-butoxyphenyl,
 - hh) 3,4-dibenzyloxyphenyl,
 - ii) 3-quinolinyl,
 - jj) 3-pyridyl,
 - kk) 4-pyridyl,
 - ll) 2,4-difluorophenyl,
 - mm) 3,4-difluorophenyl,
 - nn) 4-methylsulfinylphenyl,
 - oo) 4-methylsulfonylphenyl,
 - pp) 2-methoxyphenyl,
 - qq) 3-methoxyphenyl,
 - rr) 4-nitrophenyl,
 - ss) 4-aminomethylphenyl, and
 - tt) 2-chlorophenyl.

(R^a)_b-Ar² is selected from the group consisting of:

- 25
30
35
- a) 4-(methylthio)-phenyl,
 - b) 4-(ethylthio)-phenyl,
 - c) 3-(methylthio)-phenyl,
 - d) 2-(methylthio)-phenyl,
 - e) 3-(ethylthio)-phenyl,
 - f) 4-methylsulfonylphenyl,
 - g) 4-ethylsulfonylphenyl,
 - h) 3-methylsulfonylphenyl,
 - i) 2-methylsulfonylphenyl,
 - j) 4-methylsulfinylphenyl,
 - k) 4-ethylsulfonylphenyl,
 - l) 3-methylsulfinylphenyl,
 - m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
 - n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
 - o) 4-methoxyphenyl,

- 94 -

- 5 p) 4-hydroxyphenyl,
 q) 3-methoxyphenyl,
 r) 2-benzyloxyphenyl,
 s) 4-methylthiophen-2-yl,
 t) 4-methylthiophen-3-yl,
 u) 4-acetylamino-phenyl,
 v) 2-pyrimidinyl,
 w) phenyl,
 10 x) 4-aminomethylphenyl,
 y) 4-cyanophenyl,
 z) 4-fluorophenyl,
 aa) 4-chlorophenyl,
 bb) 4-bromophenyl,
 cc) 4-carboxyethylphenyl,
 15 dd) 2-fluorophenyl,
 ee) 3-nitrophenyl,
 ff) 4-nitrophenyl,
 gg) 3-fluorophenyl,
 hh) 4-carboxyphenyl,
 20 ii) 4-aminophenyl,
 jj) 3-aminophenyl,
 kk) 4-(O(CH₂)₃NMe₂)-phenyl,
 ll) 4-(O(CH₂)₂-piperidin-1-yl)-phenyl,
 mm) 2-methoxyphenyl,
 25 nn) 3-chlorophenyl,
 oo) 4-((4-N-COCH₃)piperazin-1-yl)-phenyl,
 pp) 4-trifluoromethylphenyl,
 qq) 4-bromothiophen-2-yl,
 rr) 5-methylthiophen-2-yl,
 30 ss) 2-benzoxazolyl,
 tt) 2-benzofuranyl,
 uu) 2,5-dimethoxyphenyl and
 vv) 4-morpholinylphenyl;
- 35 with the proviso that when (R^a)_a-Ar^I represents a), f), g), z), aa), bb), jj) or
 kk) Ar²-(R^a)_b does not represent v), w), ss) or tt);

- 95 -

(R^a)0-3-HAr is selected from the group consisting of:

- 5
- a) 4-pyridyl,
 - b) 4-(2-methylpyridyl),
 - c) 4-(2-aminopyridyl),
 - d) 4-(2-methoxypyridyl),
 - e) 4-quinolinyl,
 - f) 4-pyrimidinyl,
 - 10 g) 9-purinyl,
 - h) 7-(imidazo[4,5-b]pyridinyl),
 - i) 4-(3-methylpyridyl),
 - j) 2-pyridyl,
 - k) 3,5-dimethyl-4-pyridyl,
 - l) 3-quinolinyl,
 - 15 m) 3-pyridazinyl,
 - n) 4-(2-aminobenzyl)pyridyl, and
 - o) 4-(2-amino)pyrimidinyl;

20 R¹ is: H; and

R² is selected from the group consisting of:

- 25
- a) H,
 - b) F,
 - c) Cl,
 - d) Br,
 - e) CN,
 - f) C(O)C₁₋₆ alkyl,
 - g) C(O)C₁₋₆ alkylphenyl,
 - h) CO₂H,
 - 30 i) CO₂C₁₋₆ alkyl,
 - j) CO₂C₁₋₆ alkylphenyl,
 - k) CONH₂,
 - l) CONHC₁₋₆ alkyl,
 - m) C(O)N(C₁₋₆ alkyl)₂,
 - 35 n) SO₂NH₂,
 - o) SO₂NHC₁₋₆ alkyl,

- 96 -

- p) $\text{SO}_2\text{N}(\text{C}_{1-6} \text{ alkyl})_2$
- q) CHO and
- r) $\text{S}(\text{O})_{1-2} \text{C}_{1-6} \text{ alkyl}$.

5 9. A compound in accordance with claim 1 wherein:
 $\text{R}^a)_a\text{-Ar}^1$ is selected from the group consisting of:

- a) phenyl,
- b) 4-fluorophenyl,
- c) 4-chlorophenyl,
- 10 d) 3-fluorophenyl,
- e) 3-chlorophenyl,
- f) thiophen-2-yl,
- g) thiophen-3-yl,
- h) 4-fluorothiophen-2-yl,
- 15 i) 4-fluorothiophen-3-yl,
- j) 5-fluorothiophen-2-yl,
- k) 5-fluorothiophen-3-yl,
- l) 4-chlorothiophen-2-yl,
- m) 4-chlorothiophen-3-yl,
- 20 n) 5-chlorothiophen-2-yl,
- o) 5-chlorothiophen-3-yl,
- p) 3-methylphenyl,
- q) 3,4-dichlorophenyl,
- r) 3-hydroxyphenyl,
- 25 s) 4-hydroxyphenyl,
- t) 3,4-dihydroxyphenyl,
- u) 3-methyl-2-thiophenyl,
- v) 5-methyl-2-thiophenyl,
- w) 4-carboxymethylphenyl,
- 30 x) 3-cyanophenyl,
- y) 4-cyanophenyl,
- z) 2-pyridyl,
- aa) 2-furoyl,
- bb) 3-furoyl,
- 35 cc) 4-methylsulfinylphenyl,
- dd) 4-trifluoromethylphenyl,

- 97 -

- 5 ee) 3-trifluoromethylphenyl,
ff) 4-methylphenyl,
gg) 4-t-butoxyphenyl,
hh) 3,4-dibenzyloxyphenyl,
ii) 3-quinoliny1,
jj) 3-pyridyl,
kk) 4-pyridyl,
ll) 2,4-difluorophenyl,
mm) 3,4-difluorophenyl,
10 nn) 4-methylsulfinylphenyl,
oo) 4-methylsulfonylphenyl,
pp) 2-methoxyphenyl,
qq) 3-methoxyphenyl,
rr) 4-nitrophenyl,
15 ss) 4-aminomethylphenyl,
tt) 2-chlorophenyl,
uu) 4-thiomethyl-phenyl,
vv) 2-benzyloxy-phenyl,
ww) 4-bromophenyl,
20 xx) 4-methoxyphenyl,
yy) 4-dimethylaminophenyl,
zz) 4-((4-N-COCH₃)-piperazin-4-yl)-phenyl and
aaa) 4-(morpholinyl)-phenyl;
- 25 (Ra)_b-Ar² is selected from the group consisting of:
- 30 a) 4-(methylthio)-phenyl,
b) 4-(ethylthio)-phenyl,
c) 3-(methylthio)-phenyl,
d) 2-(methylthio)-phenyl,
e) 3-(ethylthio)-phenyl,
f) 4-methylsulfonylphenyl,
g) 4-ethylsulfonylphenyl,
h) 3-methylsulfonylphenyl,
35 i) 2-methylsulfonylphenyl,
j) 4-methylsulfinylphenyl,

- 98 -

- 5 k) 4-ethylsulfonylphenyl,
l) 3-methylsulfinylphenyl,
m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
o) 4-methoxyphenyl,
p) 4-hydroxyphenyl,
q) 3-methoxyphenyl,
r) 2-benzyloxyphenyl,
s) 4-methylthiophen-2-yl,
10 t) 4-methylthiophen-3-yl,
u) 4-acetylaminophenyl,
v) 2-pyrimidinyl,
w) phenyl,
x) 4-aminomethylphenyl,
15 y) 4-cyanophenyl,
z) 4-fluorophenyl,
aa) 4-chlorophenyl,
bb) 4-bromophenyl,
cc) 4-carboxyethylphenyl,
20 dd) 2-fluorophenyl,
ee) 3-nitrophenyl,
ff) 4-nitrophenyl,
gg) 3-fluorophenyl,
hh) 4-carboxyphenyl,
25 ii) 4-aminophenyl,
jj) 3-aminophenyl,
kk) 4-(O(CH₂)₃NMe₂)-phenyl,
ll) 4-(O(CH₂)₂-piperidin-1-yl)-phenyl,
mm) 2-methoxyphenyl,
30 nn) 3-chlorophenyl,
oo) 4-((4-N-COCH₃)piperazin-1-yl)-phenyl,
pp) 4-trifluoromethylphenyl,
qq) 4-bromothiophen-2-yl,
rr) 5-methylthiophen-2-yl,
35 ss) 2-benzoxazolyl,
tt) 2-benzofuranyl,

- 99 -

- 5 uu) 2,5-dimethoxy-phenyl and
 vv) 4-morpholinylphenyl;
 ww) 2-nitrophenyl,
 xx) 4-(CONHCH₂-2-pyridyl)-phenyl,
 yy) 4-dimethylaminophenyl,
 zz) 2-chlorophenyl,
 aaa) 4-fluoro-3-chloro-phenyl,
 aab) 4-(CONHCH₂-phenyl)-phenyl,
 aac) 4-(NHCO(CH₂)₃NMe₂)-phenyl,
 10 aad) 4-(CONHCH₂-4-pyridyl)-phenyl,
 aae) 4-(CONH(CH₂)₂-(1-piperidiny))-phenyl,
 aaf) 4-(CONH(CH₂)₂-2-(N-methylimidazolyl))-
 phenyl,
 aag) 4-(CONH(CH₂)₂-(2-pyridyl))-phenyl,
 15 aah) 4-(CONHCH₂-(3-pyridyl))-phenyl,
 aai) 3-(NHCO(CH₂)₃-NMe₂)-phenyl, and
 aaj) 4-(NHCO(CH₂)₂-(1-piperidiny))-phenyl,

20 with the proviso that when (R^a)_a-Ar¹ represents a), f), g), z), aa), bb), jj) or
 kk) Ar²-(R^a)_b does not represent v), w), ss) or tt);

(R^a)₀₋₃-HAr is selected from the group consisting of:

- 25 a) 4-pyridyl,
 b) 4-(2-methylpyridyl),
 c) 4-(2-aminopyridyl),
 d) 4-(2-methoxypyridyl),
 e) 4-quinolinyl,
 f) 4-pyrimidinyl,
 g) 9-purinyl,
 30 h) 7-(imidazo[4,5-b]pyridinyl),
 i) 4-(3-methylpyridyl),
 j) 2-pyridyl,
 k) 3,5-dimethyl-4-pyridyl,
 l) 3-quinolinyl,
 35 m) 3-pyridazinyl,
 n) 4-(2-aminobenzyl)pyridyl,

- 100 -

- 5
- o) 4-(2-amino)pyrimidinyl,
 - p) 2-quinolinyl,
 - r) 2-pyrimidinyl,
 - s) 2-pyrazinyl and
 - t) 2-(imidazo-(4,5-b)-pyridinyl;


R¹ is: substituted or unsubstituted C₁₋₁₅ alkyl; and

10 R² is selected from the group consisting of:



- 15
- a) H,
 - b) F,
 - c) Cl,
 - d) Br,
 - e) CN;
 - f) C(O)C₁₋₆ alkyl;
 - g) C(O)C₁₋₆ alkylphenyl;
 - h) CO₂H;
 - i) CO₂C₁₋₆ alkyl
 - 20 j) CO₂C₁₋₆ alkylphenyl;
 - k) CONH₂;
 - l) CONHC₁₋₆ alkyl;
 - m) C(O)N(C₁₋₆ alkyl)₂;
 - n) SO₂NH₂;
 - 25 o) SO₂NHC₁₋₆ alkyl ;
 - p) SO₂N(C₁₋₆ alkyl)₂;
 - r) CHO and
 - s) S(O)₁₋₂C₁₋₆alkyl.

30 10. A compound represented by formula I:

- 101 -




wherein:

5  and  each independently represent a 5-10 membered aryl or heteroaryl group;

a and b represents integers, 0, 1, 2 or 3, such that the sum of a plus b is 1, 2, 3 or 4;

10

 represents a heteroaryl group containing from 5 to 10 atoms, 1-3 of which are heteroatoms, 0-3 of which heteroatoms are N and 0-1 of which are O or S, said heteroaryl group being unsubstituted or substituted with 1 -3 R^a groups;

15

each R^a independently represents a member selected from the group consisting of: halo; CN, NO₂, R²¹; OR²³; SR²³; S(O)R²¹; SO₂R²¹; NR²⁰R²³; NR²⁰COR²¹; NR²⁰CO₂R²¹; NR²⁰CONR²⁰R²³; NR²⁰SO₂R²¹; NR²⁰C(NR²⁰)NHR²⁰, CO₂R²³; CONR²⁰R²³; SO₂NR²⁰R²³; SO₂NR²⁰COR²¹; SO₂NR²⁰CONR²⁰R²³; SO₂NR²⁰CO₂R²¹; OCONR²⁰R²³; OCONR²⁰SO₂R²⁰, C(O)OCH₂OC(O)R²⁰ and C(NR²⁰)NR²⁰R²³;

25 R¹ is selected from the group consisting of: H, aryl, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, and heterocyclyl, said alkyl, aryl, alkenyl, alkynyl and heterocyclyl being optionally substituted with from one to three members selected from the group consisting of: aryl, heteroaryl,

- 102 -

heterocyclyl, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{21}$, SO_2R^{21} , $SO_2N(R^{20})_2$,
 $SO_2NR^{20}COR^{21}$, $SO_2NR^{20}CON(R^{20})_2$, $NR^{20}COR^{21}$, $NR^{20}CO_2R^{21}$,
 $NR^{20}CON(R^{20})_2$, $N(R^{20})C(NR^{20})NHR^{20}$, CO_2R^{20} , $CON(R^{20})_2$,
 $CONR^{20}SO_2R^{21}$, $NR^{20}SO_2R^{21}$, $SO_2NR^{20}CO_2R^{21}$, $OCN(R^{20})_2$,
5 $CONR^{20}SO_2R^{21}$ and $CONR^{20}R^{23}$;

R^2 is selected from the group consisting of: H, C_{1-15} alkyl,
 C_{2-15} alkenyl, C_{2-15} alkynyl, halo, NO_2 and heterocyclyl, said alkyl,
alkenyl, alkynyl and heterocyclyl being optionally substituted with from one
10 to three members selected from the group consisting of: halo, heterocyclyl,
CN, aryl, heteroaryl, R^{20} , OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} ,
 $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $NR^{20}COR^{22}$,
 $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{22}C(NR^{22})NHR^{22}$, CO_2R^{20} ,
 $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$,
15 $CONR^{20}SO_2R^{22}$ and $CONR^{20}R^{23}$;

R^{20} represents a member selected from the group consisting of:
H, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and
heteroaryl, said alkyl, alkenyl and alkynyl being optionally substituted with
20 1-3 groups selected from halo, aryl and heteroaryl;

R^{21} represents a member selected from the group consisting of:
 C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl,
said alkyl, alkenyl, alkynyl, aryl and heteroaryl being optionally substituted
25 with from 1-3 of halo, heterocyclyl, aryl, heteroaryl, CN, OR^{20} ,
 $O((CH_2)_nO)_mR^{20}$, $NR^{20}((CH_2)_nO)_mR^{20}$ wherein n represents an integer of
from 2 to 4, and m represents an integer of from 1 to 3; heterocyclyl, SR^{20} ,
 $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$,
 $SO_2NR^{20}CON(R^{20})_2$, $NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$,
30 $NR^{22}C(NR^{22})NHR^{22}$, CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$,
 $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $CONR^{20}SO_2R^{22}$ $OC(O)R^{20}$,
 $C(O)OCH_2OC(O)R^{20}$ and $OCN(R^{20})_2$;

R^{22} is selected from the group consisting of: C_{1-15} alkyl, C_{2-15}
35 alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl, said alkyl,

- 103 -

alkenyl, and alkynyl being optionally substituted with 1-3 halo, aryl or heteroaryl groups;

R²³ is R²¹ or H;

5

R²⁴ is selected from COR²², CO₂R²², CON(R²⁰)₂, SO₂R²² and R²³;

and when two R²⁰ groups are present, when R²⁰ and R²¹ are present, or when R²⁰ and R²³ are present, said two R²⁰ groups, R²⁰ and R²¹ or said R²⁰ and R²³ may be taken in combination with the atoms to which they are attached and any intervening atoms and represent heterocyclyl containing from 5-10 atoms, at least one atom of which is a heteroatom selected from O, S or N, said heterocyclyl optionally containing 1-3 additional N atoms and 0-1 additional O or S atom.

15

11. A compound according to claim 10 wherein Ar¹ and Ar² are independently and optionally selected from:

- a) phenyl,
- b) pyridyl,
- 20 c) pyrimidinyl,
- d) thiophenyl,
- e) furanyl,
- f) imidazolyl,
- g) thiazolyl,
- 25 h) isothiazolyl,
- i) oxazolyl,
- j) isoxazolyl and
- k) naphthyl;

30 HAr is optionally selected from the group consisting of::

- a) pyridyl,
- b) quinolyl,
- c) purinyl,
- d) imidazolyl,
- 35 e) imidazopyridyl and
- f) pyrimidinyl;

- 104 -

- R^1 is:
- a) H or
 - b) substituted alkyl; and

5 R^2 is selected from the group consisting of::

- a) H,
- b) alkyl and
- c) halo.

10 12 A compound according to claim 10 wherein $(R^a)_a-Ar^1$ is selected from the group consisting of:

- a) phenyl,
- b) 4-fluorophenyl,
- c) 4-chlorophenyl,
- 15 d) 3-fluorophenyl,
- e) 3-chlorophenyl,
- f) thiophen-2-yl,
- g) thiophen-3-yl,
- h) 4-fluorothiophen-2-yl,
- 20 i) 4-fluorothiophen-3-yl,
- j) 5-fluorothiophen-2-yl,
- k) 5-fluorothiophen-3-yl,
- l) 4-chlorothiophen-2-yl,
- m) 4-chlorothiophen-3-yl,
- 25 n) 5-chlorothiophen-2-yl,
- o) 5-chlorothiophen-3-yl,
- p) 3-methyl phenyl,
- q) 3,4 dichlorophenyl and
- r) 3-hydroxyphenyl;

30

$Ar^2-(R^a)_b$ is selected from the group consisting of::

- a) 4-(methylthio)-phenyl,
- b) 4-(ethylthio)-phenyl,
- c) 3-(methylthio)-phenyl,
- 35 d) 2-(methylthio)-phenyl,
- e) 3-(ethylthio)-phenyl,

- 105 -

- 5 f) 4-methylsulfonylphenyl,
 g) 4-ethylsulfonylphenyl,
 h) 3-methylsulfonylphenyl,
 i) 2-methylsulfonylphenyl,
 j) 4-methylsulfinylphenyl,
 k) 4-ethylsulfonylphenyl,
 l) 3-methylsulfinylphenyl,
 m) 4-(N-methyl-N-benzyl)aminomethylphenyl,
 n) 3-(N-methyl-N-benzyl)aminomethylphenyl,
 10 o) 4-methoxyphenyl,
 p) 4-hydroxyphenyl,
 q) 3-methoxyphenyl,
 r) 2-benzyloxyphenyl,
 s) 4-methylthiophen-2-yl,
 15 t) 4-methylthiophen-3-yl,
 u) 4-acetylamino-phenyl and
 v) 2-pyrimidinyl;

HAr is selected from the group consisting of::

- 20 a) 4-pyridyl,
 b) 4-(2-methylpyridyl),
 c) 4-(2-aminopyridyl),
 d) 4-(2-methoxypyridyl),
 e) 4-quinolinyl,
 25 f) 4-pyrimidinyl,
 g) 9-purinyl,
 h) 7-(imidazo[4,5-b]pyridinyl), and
 i) 4-(3-methylpyridyl)

30 R¹ is: H and


R² is selected from the group consisting of:

- a) H,
 b) F,
 35 c) Cl and
 d) Br.

13. A compound in accordance with claim 1 selected from

Table IV:

5

TABLE IV				
				
Ex #	$(R^a)_a$ -Ar ¹	Ar ² -(R ^a) _b	$(R^a)_{0-3}$ -HAr	FAB ms
2	Ph-4-F	Ph-4-OMe	4-Pyridyl	345
3	Ph-4-F	Ph-2,5-di-OMe	4-Pyridyl	375
4	Ph-4-F	Ph-4-Br	4-Pyridyl	393/395
5	Ph-4-F	Ph-4-Cl	4-Pyridyl	349
6	Ph-4-F	Ph-4-OMe	2-Pyridyl	345
7	Ph-4-F	Ph-4-Br	2-Pyridyl	393/395
8	Ph-4-F	Ph-4-Cl	2-Pyridyl	349
9	Ph-4-F	Ph-2,5-di-OMe	2-Pyridyl	375
10	Ph	Ph-4-OMe	4-Pyridyl	327
11	Ph	Ph-4-Cl	4-Pyridyl	331
12	Ph	Ph-2,5-di-OMe	4-Pyridyl	357
13	Ph	Ph-4-F	4-Pyridyl	315
14	Ph-4-Cl	Ph	4-Pyridyl	331
15	Ph-4-F	Ph-4-SMe	4-Pyridyl	361
16	Ph-4-SMe	Ph	4-Pyridyl	325
17	4-CF ₃ -Ph	Ph	4-Pyridyl	365

- 107 -

18	4-Me-Ph	Ph	4-Pyridyl	311
19	4-tBuO-Ph	Ph	4-Pyridyl	369
20	3,4-Cl-Ph	Ph	4-Pyridyl	366
21	3,4-di-(OBn)-Ph	Ph	4-Pyridyl	509
22	3-Cl-Ph	Ph	4-pyridyl	331
23	4-CN-Ph	Ph	4-Pyridyl	322
24	4-(COOMe)-Ph	Ph	4-Pyridyl	355
25	Ph	4-CO ₂ Et-Ph	4-Pyridyl	369
26	Ph	2-F-Ph	4-Pyridyl	315
27	Ph	3-NO ₂ -Ph	4-Pyridyl	342
28	5-Me-2-thiophenyl	Ph	4-Pyridyl	317
29	3-Me-2-thiophenyl	Ph	4-Pyridyl	317
30	4-F-Ph	4-SMe-Ph	4-(3-Me)-Pyridyl	375
31	Ph	2-NO ₂ -Ph	4-Pyridyl	342
32	Ph	3-F-Ph	4-Pyridyl	315
33	Ph	4-NO ₂ -Ph	4-Pyridyl	342
34	4-F-Ph	Ph	4-Pyridyl	315
35	2,4-F-Ph	Ph	4-Pyridyl	333
36	3-CN-Ph	Ph	4-Pyridyl	322
37	3,4-F-Ph	Ph	4-Pyridyl	333
38	4-F-Ph	4-(SMe)-Ph	4-quinolinyl	411
39	2-Cl-Ph	Ph	4-pyridyl	331
40	3-Cl-Ph	4-(SMe)-Ph	4-pyridyl	377
41	4-F-Ph	Ph	4-pyrimidinyl	316
42	4-F-Ph	4-(SMe)-Ph	4-(2-methyl)-pyridyl	375
43	3,4-di-F-Ph	4-(SMe)-Ph	4-pyridyl	379.


14. A compound in accordance with claim 1 selected from

Table V:



5 :

TABLE V





- 108 -

			
Example #	Aryl ¹	Har	FAB ms
45	4-F-phenyl	4-(3-Me)-pyridyl	391
46	4-F-phenyl	4-quinolinyl	427
47	3-Cl-phenyl	4-pyridyl	393
48	4-F-phenyl	4-(2-Me)-pyridyl	391
49	3,4-F-phenyl	4-pyridyl	395.




15. A compound in accordance with claim 1 selected from Table VI:

TABLE VI		
		
Example #	Aryl ² =	FAB ms
55	4-(CONHCH ₂ -phenyl)-phenyl 	430

- 109 -

56	<p>4-(NHCO(CH₂)₂-(1-piperidinyl))-phenyl</p> 	451
57	<p>4-(CONHCH₂-4-pyr)-phenyl</p> 	431
58	<p>4-(CONH(CH₂)₂-(1-piperidinyl))-phenyl</p> 	451
59	<p>4-(CONH(CH₂)₂-2-(N-methylimidazolyl)phenyl</p> 	447

- 110 -


60	<p>4-(CONH(CH₂)₂-(2-pyridyl))-phenyl</p> 	445
61	<p>4-(CONHCH₂-(3-pyridyl))-phenyl</p> 	431
62	<p>3-(NHCO(CH₂)₃-NMe₂)-phenyl</p> 	425

16. A compound in accordance with claim 1 which is:
- 2-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-3-(4-pyridyl) pyrrole;
 - 5 2-(3-chlorophenyl)-5-(4-methylsulfonylphenyl)-3-(4-pyridyl) pyrrole;
 - 2,5-bis-(4-fluorophenyl)-3-(4-pyridyl)-pyrrole;
 - 2-(4-fluorophenyl)-5-(4-methylsulfinylphenyl)-3-(4-pyridyl)-pyrrole;
 - 2-(4-methylsulfinyl phenyl)-5-(phenyl)-3-(4-pyridyl)-pyrrole;
 - 2-(4-phenyl)-5-(4-carboxyphenyl)-3-(4-pyridyl)-pyrrole;
 - 10 2-(4-phenyl)-5-(4-aminophenyl)-3-(4-pyridyl)-pyrrole;
 - 2-(4-phenyl)-5-(3-aminophenyl)-3-(4-pyridyl)-pyrrole;

- 111 -

2-(4-phenyl)-5-(4-CONHCH₂-2-pyridyl)-phenyl-3-(4-pyridyl)-pyrrole;
 2-(4-hydroxyphenyl)-5-(phenyl)-3-(4-pyridyl)-pyrrole or
 2-(3,4-(OH)₂-phenyl)-5-(phenyl)-3-(4-pyridyl)-pyrrole.

- 5 17. A compound in accordance with claim 1 selected from Table VII:

TABLE VII				
				
Ex#	(R ^a) _a -Ar ¹	Ar ² -(R ^a) _b	(R ^a) ₀₋₃ -HAr	R ²
67	Ph	Ph-4-F	3-quinolinyl	H
68	Ph	Ph-4-F	4-pyrimidinyl	H
69	Ph	Ph-4-F	3-pyridazinyl	H
70	Ph	Ph-4-F	2-pyrazinyl	H
71	Ph	Ph-4-CN	4-pyridyl	H
72	Ph	Ph-2-OMe	4-pyridyl	H
73	Ph	Ph-3-OMe	4-pyridyl	H
74	Ph-3,4-di-F	Ph-4-S(O)-Me	4-pyridyl	H
75	Ph	Ph-4-NMe ₂	4-pyridyl	H
76	Ph	4-(4-(N-COCH ₃) piperazinyl)-Ph	4-pyridyl	H
77	Ph	4-(morpholinyl)-Ph	4-pyridyl	H
78	Ph	Ph-2-Cl	4-pyridyl	H
79	Ph	Ph-3-Cl	4-pyridyl	H
80	Ph	Ph-4-CF ₃	4-pyridyl	H
81	Ph	Ph-4-S-Me	4-pyridyl	H
82	Ph	Ph-4-S(O)-Me	4-pyridyl	H


- 112 -

83	Ph-4-F	(4-methyl)-2-thiophenyl	4-pyridyl	H
84	Ph-4-F	(4-bromo)-2-thiophenyl	4-pyridyl	H
85	Ph-4-F	Ph-4-F-3-Cl	4-pyridyl	H
86	Ph-4-F	2-benzoxazolyl	4-pyridyl	H
87	Ph-4-F	2-benzofuranyl	4-pyridyl	H
88	Ph-4-F	4-(O(CH ₂) ₃ NMe ₂)-Ph	4-pyridyl	H
89	Ph-4-F	4-(O(CH ₂) ₂ -piperidin-1-yl)-Ph	4-pyridyl	H
90	Ph-4-Cl	Ph-4-F	4-pyridyl	H
91	Ph-3-Cl	Ph-4-F	4-pyridyl	H
92	Ph-2-Cl	Ph-4-F	4-pyridyl	H
93	Ph-3,4-di-Cl	Ph-4-F	4-pyridyl	H
94	Ph-3-CF ₃	Ph-4-F	4-pyridyl	H
95	Ph-4-S-Me	Ph-4-F	4-pyridyl	H
96	Ph-4-S(O)-Me	Ph-4-F	4-pyridyl	H
97	Ph-2-OBn	Ph-4-F	4-pyridyl	H
98	Ph-4-Br	Ph-4-F	4-pyridyl	H
99	Ph-2-OMe	Ph-4-F	4-pyridyl	H
100	Ph-3-OMe	Ph-4-F	4-pyridyl	H
101	Ph-4-OMe	Ph-4-F	4-pyridyl	H
102	Ph-4-NO ₂	Ph-4-F	4-pyridyl	H
103	Ph-4-NMe ₂	Ph-4-F	4-pyridyl	H
104	Ph-4-(4-N-COCH ₃)-piperazinyl	Ph-4-F	4-pyridyl	H
105	Ph-4-morpholinyl	Ph-4-F	4-pyridyl	H
106	2-thiophenyl	Ph-4-F	4-pyridyl	H
107	3-thiophenyl	Ph-4-F	4-pyridyl	H
108	2-furoyl	Ph-4-F	4-pyridyl	H
109	3-furoyl	Ph-4-F	4-pyridyl	H
110	Ph-3-Cl	Ph-4-S(O)Me	4-(2-Me)-pyridyl	H
111	Ph-3,4-di-F	Ph-4-S(O)Me	4-(2-Me)-pyridyl	H
112	Ph-3,4-di-F	Ph-4-S(O)Me	4-pyridyl	F
113	Ph-3-CF ₃	Ph	4-pyridyl	H
114	Ph-4-F	Ph-4-S(O)Me	4-(2-amino benzyl)pyridyl	H
115	3-quinolinyl	Ph	4-pyridyl	H









- 113 -

116	Ph-4-F	4-(SMe)-Ph	4-pyridyl	Br
117	Ph-3,4-di-F	Ph	4-pyridyl	CO ₂ Et
118	Ph-3,4-di-F	Ph	4-pyridyl	CN
119	3-(CF ₃)-Ph	Ph	4-pyridyl	CO ₂ Et
120	3-(CF ₃)-Ph	Ph	4-pyridyl	CN
121	Ph	Ph-4-F	3-methyl-4-pyridyl	H
122	Ph-3-Cl	Ph-4-S(O)Me	4-(2-amino)pyrimidinyl	H
123	Ph	Ph-4-F	3,5-dimethyl-4-pyridyl	H
Bn = benzyl, Ph = phenyl, Me = methyl Et = ethyl .				







18. A compound in accordance with claim 1 represented by the structural formula:



- 114 -


 <chem>Nc1ccc(cc1)c2c(c[nH]2)c3ccccc3c4ccncc4</chem>	 <chem>CCC(=O)c1c(c[nH]1c2ccc(Br)cc2)c3ccccc3c4cc(F)cc(F)c4c5ccncc5</chem>
 <chem>Oc1ccc(cc1)c2c(c[nH]2)c3ccccc3c4ccncc4</chem>	 <chem>Oc1cc(O)ccc1c2c(c[nH]2)c3ccccc3c4ccncc4</chem>
 <chem>CCOC(=O)c1c(c[nH]1c2ccccc2)c3cc(F)cc(F)c3c4ccncc4</chem>	 <chem>c1ccccc1COCC(=O)c2c(c[nH]2c3cc(F)cc(F)c3)c4ccncc4</chem>
 <chem>OC(=O)c1c(c[nH]1c2ccccc2)c3cc(F)cc(F)c3c4ccncc4</chem>	 <chem>CN(C)C(=O)c1c(c[nH]1c2ccccc2)c3cc(F)cc(F)c3c4ccncc4</chem>

- 115 -



- 116 -

19. A compound in accordance with claim 1 selected from Table VIII:

TABLE VIII				
				
Ex.	$(R^a)_a$ —Ar ¹	Ar ² — $(R^a)_b$	$(R^a)_{0-3}$ —HAr	R ²
134	Ph-4-F	Ph-4-OMe	4-pyridyl	C(O)Me
135	Ph-4-F	Ph-2,5-di-OMe	4-pyridyl	C(O)Me
136	Ph-4-F	Ph-4-Br	4-pyridyl	C(O)Propyl
137	Ph-4-F	Ph-4-Cl	4-pyridyl	C(O)Ethyl
138	Ph-4-F	Ph-4-OMe	2-pyridyl	C(O)Me
139	Ph-4-F	Ph-4-Br	2-pyridyl	C(O)Me
140	Ph-4-F	Ph-2,5-di-OMe	2-pyridyl	C(O)Me
141	Ph	Ph-4-OMe	4-pyridyl	C(O)Me
142	Ph	Ph-4-Cl	4-pyridyl	C(O)Me
143	Ph	Ph-2,5-di-OMe	4-pyridyl	C(O)Me
144	Ph	Ph-4-F	4-pyridyl	C(O)Me
145	Ph-4-Cl	Ph	4-pyridyl	C(O)Me
146	Ph-4-F	Ph-4-SMe	4-pyridyl	C(O)Me
147	Ph-4-SMe	Ph	3-methyl-4-pyridyl	C(O)Me
148	Ph-4-F	Ph-4-SMe	3-methyl-4-pyridyl	CN
149	Ph-4-F	Ph-4-S(O)Me	3-methyl-4-pyridyl	CN
150	Ph-3-Cl	Ph-4-Cl	4-quinolyl	CN
151	Ph-4-F	Ph-4-Cl	2-methyl-4-pyridyl	CN

- 117 -


152	Ph	Ph-4-F	3,5-dimethyl-4-pyridyl	CN
153	Ph	Ph-4-F	3-quinoliny	CN
154	Ph	Ph-4-F	4-quinoliny	CN
155	Ph	Ph-4-F	2-quinoliny	CN
156	Ph	Ph-4-F	2-pyrimidiny	CN
157	Ph	Ph-4-F	4-pyrimidiny	CN
158	Ph	Ph-4-F	3-pyridaziny	CN
159	Ph	Ph-4-F	2-pyraziny	CN
160	Ph	Ph-4-F	2-pyrimidiny	CN
161	Ph	Ph-4-F	4-pyrimidiny	CN
162	Ph	Ph-4-F	2-imidazo-(4,5-b)-pyridiny	CN
163	Ph	Ph-4-F	7-imidazo-(4,3-b)-pyridiny	CN
164	Ph	Ph-4-F	4-pyridyl	COMe
165	Ph	Ph-4-F	4-pyridyl	SO ₂ Me
166	Ph	Ph-4-CN	4-pyridyl	COMe
167	Ph	Ph-2-OMe	4-pyridyl	COMe
168	Ph	Ph-3-OMe	4-pyridyl	CN
169	Ph	Ph-4-OMe	4-pyridyl	CO ₂ Et
170	Ph	Ph-4-NO ₂	4-pyridyl	CN
171	Ph	Ph-4-NMe ₂	4-pyridyl	CN
172	Ph	4-(4-(N-COCH ₃)-piperaziny)-Ph	4-pyridyl	CN
173	Ph	4-(morpholiny)-Ph	4-pyridyl	CN
174	Ph	Ph-2-Cl	4-pyridyl	CN
175	Ph	Ph-3-Cl	4-pyridyl	C(O)Me
176	Ph	Ph-4-CF ₃	4-pyridyl	SO ₂ Me
177	Ph	Ph-4-S-Me	4-pyridyl	CN
178	Ph	Ph-4-S(O)-Me	4-pyridyl	CN

- 118 -

179	Ph-4-F	(4-methyl) thiophen-2yl	4-pyridyl	CN
180	Ph-4-F	(3-methyl) thiophen-2yl	4-pyridyl	C(O)Me
181	Ph-4-F	(4-bromo) thiophen-2yl	4-pyridyl	SO ₂ Me
182	Ph-4-F	(5-methyl) thiophen-2yl	4-pyridyl	CN
183	Ph-4-F	Ph-4-F-3-Cl	4-pyridyl	CN
184	Ph-4-F	2-benzoxazolyl	4-pyridyl	CN
185	Ph-4-F	2-benzofuranyl	4-pyridyl	CN
186	Ph-4-F	4-(O(CH ₂) ₃ - NMe ₂)-Ph	4-pyridyl	CN
187	Ph-4-F	4-(O(CH ₂) ₂ N- piperidiny)-Ph	4-pyridyl	CN
188	Ph-4-Cl	Ph-4-F	4-pyridyl	CN
189	Ph-3-Cl	Ph-4-F	4-pyridyl	CN
190	Ph-2-Cl	Ph-4-F	4-pyridyl	CN
191	Ph-3,4-di-Cl	Ph-4-F	4-pyridyl	CN
192	Ph-3-CF ₃	Ph-4-F	4-pyridyl	CN
193	Ph-4-S-Me	Ph-4-F	4-pyridyl	CN
194	Ph-4-S(O)-Me	Ph-4-F	4-pyridyl	CN
195	Ph-2-OBn	Ph-4-F	4-pyridyl	CN
196	Ph-4-Br	Ph-4-F	4-pyridyl	CN
197	Ph-2-OMe	Ph-4-F	4-pyridyl	CN
198	Ph-3-OMe	Ph-4-F	4-pyridyl	CN
199	Ph-4-OMe	Ph-4-F	4-pyridyl	CN
200	Ph-4-NO ₂	Ph-4-F	4-pyridyl	CN
201	Ph-4-NMe ₂	Ph-4-F	4-pyridyl	CN
202	4-(4-(N-COCH ₃)- piperazinyl)-Ph	Ph-4-F	4-pyridyl	CN
203	4-(morpholinyl)-Ph	Ph-4-F	4-pyridyl	CN
Me = methyl, Et = ethyl, Ph = phenyl, Bn = benzyl,				

- 119 -

20. A compound in accordance with claim 1 falling within the table IX:

TABLE IX				
				
R ¹	(R ^a) _a -Ar ¹	Ar ² -(R ^a) _b	(R ^a) ₀₋₃ -HAr	R ²
H	4-F-Ph	4-F-Ph	4-Pyr	H
H	4-F-Ph	4-OMe-Ph	4-Pyr	H
H	4-F-Ph	2,5-di-(OMe)-Ph	4-Pyr	H
H	4-F-Ph	4-Br-Ph	4-Pyr	H
H	4-F-Ph	4-Cl-Ph	4-Pyr	H
H	Ph	4-OMe-Ph	4-Pyr	H
H	Ph	4-Cl-Ph	4-Pyr	H
H	Ph	2,5-di-(OMe)-Ph	4-Pyr	H
H	Ph	4-F-Ph	4-Pyr	H
H	4-F-Ph	4-SMe-Ph	4-Pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-Pyr	H
H	4-Cl-Ph	Ph	4-Pyr	H
H	4-SMe-Ph	Ph	4-Pyr	H
H	4-S(O)Me-Ph-4-Pyr	Ph	H	Ph
H	4-CF ₃ -Ph	Ph	4-Pyr	H
H	4-Me-Ph	Ph	4-Pyr	H
H	4-OH-Ph	Ph	4-Pyr	H
H	3,4-di-Cl-Ph	Ph	4-Pyr	H

- 120 -

H	3,4-di-OH-Ph	Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	CO ₂ Et
H	3-Cl-Ph	Ph	4-pyr	H
H	4-CN-Ph	Ph	4-Pyr	H
H	Ph	4-CO ₂ Et-Ph	4-Pyr	H
H	Ph	2-F-Ph	4-Pyr	H
H	Ph	3-NO ₂ -Ph	4-Pyr	H
H	5-Me-thiophen-2-yl	Ph	4-Pyr	H
H	3-Me-thiophen-2-yl	Ph	4-Pyr	H
H	3-quinolinyl	Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	CN
H	4-F-Ph	4-(SMe)-Ph	4-(3-Me)-Pyr	H
H	4-F-Ph	4-(SMe)-Ph	4-(3-Me)-Pyr	H
H	Ph	2-(NO ₂)-Ph	4-Pyr	H
H	Ph	3-F-Ph	4-Pyr	H
H	Ph	4-CO ₂ H-Ph	4-Pyr	H
H	Ph	3-NH ₂ -Ph	4-Pyr	H
H	Ph	4-NO ₂ -Ph	4-Pyr	H
H	4-F-Ph	Ph	4-Pyr	H
H	2,4-di-F-Ph	Ph	4-Pyr	H
H	3-CN-Ph	Ph	4-Pyr	H
H	3,4-di-F-Ph	Ph	4-Pyr	H
H	Ph	4-NH ₂ -Ph	4-Pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-quinolinyl	H
H	4-F-Ph	4-SMe-Ph	4-pyr	Br
H	4-F-Ph	4-SMe-Ph	4-quinolinyl	H
H	Ph	4-(CONHCH ₂ -phenyl)-Ph	4-pyr	H
H	Ph	4-(NHCO(CH ₂) ₃ -NMe ₂)-Ph	4-pyr	H
H	Ph	4-(CONHCH ₂ -4-pyr)-Ph	4-pyr	H
H	Ph	4-(CONH(CH ₂) ₂ -(1-piperidinyl))-Ph	4-pyr	H
H	Ph	4-(CONH(CH ₂) ₂ -2-(N-methylimidazolyl))-Ph	4-pyr	H
H	Ph	4-(CONHCH ₂ -(2-pyridyl))-Ph	4-pyr	H

- 121 -

H	Ph	4-(CONH(CH ₂) ₂ -(2-pyridyl))-Ph	4-pyr	H
H	Ph	4-(CONHCH ₂ -(3-pyridyl))-Ph	4-pyr	H
H	Ph	3-(NHCO(CH ₂) ₃ NMe ₂)-Ph	4-pyr	H
H	Ph	4-(NHCO(CH ₂) ₂ -(1-piperidyl))-Ph	4-pyr	H
H	2-Cl-Ph	Ph	4-pyr	H
H	3-Cl-Ph	4-SMe-Ph	4-pyr	H
H	4-F-Ph	Ph	4-pyrimidinyl	F
H	4-F-Ph	4-SMe-Ph	4-(2-methyl)-pyr	H
H	3-Cl-Ph	4-S(O)Me-Ph	4-pyr	H
H	4-F-Ph	4-S(O)Me-Ph	4-(2-methyl)-pyr	H
H	3,4-di-F-Ph	4-SMe-Ph	4-pyr	H
H	3,4-di-F-Ph	4-S(O)Me-Ph	4-Pyr	H
H	4-F-Ph	4-SO ₂ Me-Ph	4-pyr	H
H	3,4-di-F-Ph	Ph	4-pyr	CO ₂ Et
H	3,4-di-F-Ph	Ph	4-pyr	CN
H	3-CF ₃ -Ph	Ph	4-pyr	CO ₂ Et
H	3-CF ₃ -Ph	Ph	4-pyr	CN
H	3-Cl-Ph	4-SO ₂ Me-Ph	4-pyr	H

21. A method of treating a cytokine mediated disease in a mammal in need of such treatment, which comprises administering to said mammal an effective cytokine interfering amount of a compound of claim 1.

22. The method according to claim 21 wherein the cytokine inhibited is IL-1.

23. The method according to claim 21 wherein the cytokine inhibited is IL-6.

24. The method according to claim 21 wherein the cytokine inhibited is TNF.

- 122 -

25. The method according to claim 21 wherein the cytokine inhibited is IL-8.

5 26. The method according to claim 21 wherein the cytokine mediated disease is septic shock, endotoxic shock, gram negative sepsis or toxic shock syndrome.

10 27. The method according to claim 21 wherein the cytokine mediated disease is bone resorption disease, graft versus host reaction, atherosclerosis, arthritis, osteoarthritis, rheumatoid arthritis, gout, psoriasis or a topical inflammatory disease state.

15 28. The method according to claim 21 wherein the cytokine mediated disease is adult respiratory distress syndrome, asthma or chronic pulmonary inflammatory disease.

20 29. The method according to claim 21 wherein the cytokine mediated disease is cardiac or renal reperfusion injury, thrombosis or glomerulonephritis.

30 30. The method according to claim 21 wherein the cytokine mediated disease is Crohn's disease, ulcerative colitis or inflammatory bowel disease.

25 31. The method according to claim 21 wherein the cytokine mediated disease is cachexia.

30 32. The method according to claim 21 wherein the cytokine mediated disease is a viral infection.

35 33. A method of treating inflammation mediated by excess production of prostaglandins in a human in need of such treatment, which comprises administering to said human an effective cytokine interfering amount of a compound of claim 1.

- 123 -

34. A pharmaceutical composition comprising a compound according to claim 1 in combination with a pharmaceutically acceptable carrier.

5 35. A pharmaceutical composition made by combining a compound according to claim 1 and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12922

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE file Registry, CAPlus

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SILVESTRI et al. Electrochemical Oxidation of 2,4,5-triaryl-substituted pyrroles. II. Oxidation Dimerization of 4,5-Diphenyl-2-Mesitylpyrrole. J. Het. Chem. March-April 1989, Vol. 26, No. 2, pages 489-492.	1-20
A	SILVESTRI et al. Electrochemical Oxidation of Substituted Pyrroles. IV. The Influence of Soluble Bases. J. Het. Chem. December 1992, Vol. 29, No. 7, pages 1847-1850.	1-20
X	KONAKAHARA et al. One-pot'synthesis of pyrroles from N-silyl-1-azaallyl anions. Heterocycles. 01 May 1993. Vol. 35, No. 2, pages 1171-1184, especially page 1175.	1-12
----		----
A		13-20

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance		
"E" earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

04 NOVEMBER 1996

Date of mailing of the international search report

19 NOV 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RICHARD S. MYERS, JR.

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US96/12922**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PETRUSO et al. Oxidative reactions of polyarylpyrroles. Part 3. Oxidation Trimerization of 2,5-Diphenylpyrrole. J.C.S. Perkins I. October 1981, Vol. 10, pages 2642-2646.	1-20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/12922

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/44, 31/50, 31/445, 31/495, 31/505, 31/535; C07D 211/68, 211/80, 213/02, 239/02, 401/00, 401/04, 471/02, 473/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/235.5, 252, 253, 256, 303, 274, 275, 314, 318, 341, 343; 544/124, 238, 272.2, 318, 331, 332, 335, 363, 364, 405; 546/119, 167, 194, 208, 256, 275.1, 276.4

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/235.5, 252, 253, 256, 303, 274, 275, 314, 318, 341, 343; 544/124, 238, 272.2, 318, 331, 332, 335, 363, 364, 405; 546/119, 167, 194, 208, 256, 275.1, 276.4